

VEGETATION-ENVIRONMENT RELATIONSHIPS IN THE
PRAIRIE-FOREST TRANSITION ZONE IN MINNESOTA

A THESIS

SUBMITTED TO THE FACULTY OF THE GRADUATE SCHOOL
OF THE UNIVERSITY OF MINNESOTA

Part II

2

Part 2

By

LARRY ALBERT DREW

IN PARTIAL FULFILLMENT OF THE REQUIREMENTS
FOR THE DEGREE OF
DOCTOR OF PHILOSOPHY

Degree Granted
December, 1973

4.0 LITERATURE CITED

- Ackoff, R.L. 1962. Scientific method. Optimizing applied research decisions. John Wiley and Sons, Inc., New York. 464p.
- Adams, C.C. 1902. Southeastern United States as a center of geographical distribution of flora and fauna. Biol. Bull. 3: 115-131.
- Adams, C.C. 1905. The postglacial dispersal of the North American biota. Biol. Bull. 9: 53-71.
- Adams, W.A. 1973. The effect of organic matter on the bulk and true densities of some uncultivated podzolic soils. J. Soil Sci. 24(1): 10-17.
- Agr. Exp. Sta. of Illinois, Indiana, Iowa, Kansas, Michigan, Minnesota, Missouri, Nebraska, North Dakota, Ohio, South Dakota, and Wisconsin and the U.S. Dept. Agr. 1960. Soils of the North Central region of the United States. Agr. Exp. Sta., Univ. of Wisconsin, Madison. 192p.
- Ahlmann, H.W. 1952. Glacier variations and climatic fluctuations. Bowman Mem. Lect., Ser. 3, Amer. Geogr. Soc., New York. 51p.
- Aikman, J.M. 1926. Distribution and structure of the forests of eastern Nebraska. Nebraska Univ., Univ. Stud. 26: 1-95.
- /Aikman, J.M. 1928. Competition studies in the ecotone between prairie and woodland. Proc. Iowa Acad. Sci. 35: 99-103.
- Aikman, J.M. and A.W. Smelser. 1938. The structure and environment of forest communities in central Iowa. Ecology 19: 141-150.
- /Albertson, F.W. and J.E. Weaver. 1945. Injury and death or recovery of trees in prairie climate. Ecol. Monogr. 15: 393-433.
- /Aldous, A.E. 1934. Effects of burning on Kansas bluestem pastures. Kansas Agr. Exp. Sta. Bull. 38. 65p.
- /Allen, J.A. 1870. The flora of the prairies. Amer. Natur. 4: 577-585.
- /Allen, J.A. 1871. The fauna of the prairies. Amer. Natur. 5: 4-9.

American Chemical Society. 1969. Cleaning our environment. The chemical basis for action. Amer. Chem. Soc., Washington, D.C. 249p.

Anderson, M.C. 1964. Light relations of terrestrial plant communities and their measurement. Biol. Rev. 39: 425-486.

Anderson, R.C. 1973. The use of fire as a management tool on the Curtis Prairie. Proc. Ann. Tall Timbers Fire Ecol. Conf. 12: 23-25.

Anderson, R.Y. 1961. Solar-terrestrial climatic patterns in varved sediments. New York Acad. Sci., Ann. 95, Art. 1: 424-439.

Anonymous. 1838. Prairies of Ohio. Amer. J. Sci., Arts 33: 230-236.

Arneman, H.F. 1963. Soils of Minnesota. Univ. Minnesota Agr. Ext. Bull. 278. 8p.

Arneman, H.F., A.D. Khan, and P.R. McMiller. 1958. Physical, chemical, and mineralogical properties of related Minnesota prairie soils. Univ. Minnesota Agr. Exp. Sta. Tech. Bull. 227. 47p.

Arneman, H.F. and D. Pluth. 1963. Field forest soils. Unpubl. Lab. Manual. Dept. Soils, Univ. Minnesota, St. Paul. 42p.

Atwater, C. 1819. On the prairies and barrens of the west. Amer. J. Sci., more esp. of Mineralogy, Geology, and the other branches of Natural History; including also Agriculture and the ornamental as well as useful Arts 1 (2nd ed., 1st Ser.): 116-125.

Ayres, H.B. 1892. IV. The woods of Minnesota, pp. 128-137. In, The Geological and Natural History Survey of Minnesota, 19th Ann. Rep., 1890, N.H. Winchell, ed. 255p.

Baker, D.G. and D.A. Haines. 1969. Solar radiation and sunshine duration relationships in the North-Central Region and Alaska. Univ. Minnesota Agr. Exp. Sta. Tech. Bull. 262. (North Central Regional Pub. 195). 372p.

Baker, D.G., D.A. Haines, and J.H. Strub, Jr. 1967. Climate of Minnesota. Part V. Precipitation facts, normals, and extremes. Univ. Minnesota Agr. Exp. Sta. Tech. Bull. 254. 44p.

- Baker, D.G. and J.H. Strub, Jr. 1963a. Climate of Minnesota. Part I. Probability of occurrence in the spring and fall of selected low temperatures. Univ. Minnesota Agr. Exp. Sta. Tech. Bull. 243. 40p.
- Baker, D.G. and J.H. Strub, Jr. 1963b. Climate of Minnesota. Part II. The agricultural and minimum-temperature-free seasons. Univ. Minnesota Agr. Exp. Sta. Tech. Bull. 245. 32p.
- Baker, D.G. and J.H. Strub, Jr. 1965. Climate of Minnesota. Part III. Temperature and its application. Univ. Minnesota Agr. Exp. Sta. Tech. Bull. 248. 64p.
- Baker, D.G. and J.B. Swan. 1966. Climate of Minnesota. Part IV. Spring soil temperatures. Univ. Minnesota Agr. Exp. Sta. Misc. Rep. 67. 11p.
- Bakuzis, E.V. 1959. Synecological coordinates in forest classification and in reproduction studies. Ph.D. Thesis, Univ. Minnesota. 244p.
- Bakuzis, E.V. 1960. Structural organization of forest ecosystems. Proc. Minnesota Acad. Sci. 27: 97-103.
- Bakuzis, E.V. 1962. Synecological coordinates and investigations of forest ecosystems. Proc. 13th Congr. Int. Union Forest. Res. Organ. (Vienna, 1961) 2(1), 11p.
- Bakuzis, E.V. 1966. Provisional list of synecological coordinates of Minnesota forest plant species. Unpubl. Mimeo, Coll. Forest., Univ. Minnesota, St. Paul. 11p.
- Bakuzis, E.V. 1967. Some characteristics of forest ecosystem space in Minnesota. Paper 14th Congr. Int. Union Forest. Res. Organ. (Munich) 2: 107-125.
- Bakuzis, E.V. 1969a. Ecosystem coordinates and forest ecosystems. Summary of the paper presented at the XI Int. Bot. Congr. (Seattle, Washington). Mimeo. 19p.
- Bakuzis, E.V. 1969b. Forestry viewed in an ecosystem perspective, pp. 189-258. In, The ecosystem concept in natural resource management, G.M. VanDyne, ed. Academic Press, New York. 383p.
- Bakuzis, E.V., D.P. Duncan, and H.L. Hansen. 1960a. Application of the method of synecological coordinates in a study of forest geographical subdivisions of Minnesota. Minnesota Forest. Notes No. 99. 2p.

- ✓ Bakuzis, E.V., D.P. Duncan, and H.L. Hansen. 1960b. Cover relationships in Minnesota forest communities. Minnesota Forest. Notes No. 100. 2p.
- Bakuzis, E.V., D.P. Duncan, and H.L. Hansen. 1960c. Suggestions for further development of the forest cover type system of eastern North America. Minnesota Forest. Notes No. 92. 2p.
- Bakuzis, E.V. and H.L. Hansen. 1959. A provisional assessment of species synecological requirements in Minnesota forests. Minnesota Forest. Notes No. 84. 2p.
- Bakuzis, E.V. and H.L. Hansen. 1960a. Some relationships between vegetational indication and broad soil groups in Minnesota forest communities. Minnesota Forest. Notes No. 90. 2p.
- Bakuzis, E.V. and H.L. Hansen. 1960b. Use of ecographs in analyzing species-environmental relationships in forest communities. Minnesota Forest. Notes No. 91. 2p.
- Bakuzis, E.V. and H.L. Hansen. 1962a. Distribution of balsam fir reproduction and basal area in the edaphic field of forest communities in the central pine section of Minnesota. Minnesota Forest. Notes No. 120. 2p.
- Bakuzis, E.V. and H.L. Hansen. 1962b. Ecographs of herb species of Minnesota forest communities. Minnesota Forest. Notes No. 118. 2p.
- Bakuzis, E.V. and H.L. Hansen. 1962c. Ecographs of shrubs and other undergrowth species of Minnesota forest communities. Minnesota Forest. Notes No. 117. 2p.
- Bakuzis, E.V., H.L. Hansen, and D.P. Duncan. 1960. Some characteristics of geographic and synecological distribution of balsam fir (Abies balsamea (L.) Mill.) in Minnesota. Minnesota Forest. Notes No. 101. 2p.
- Bakuzis, E.V., H.L. Hansen, and V. Kurmis. 1962. Weighted site indices in relation to some characteristics of the edaphic field of the central pine section of Minnesota forests. Minnesota Forest. Notes No. 119. 2p.
- Bard, G.E. 1949. The mineral nutrient content of the annual parts of herbaceous species growing on three New York soil types varying in limestone content. Ecology 30: 384-389.

- Baumgartner, A. 1965. The heat, water and carbon dioxide budget of plant cover: Methods and measurements, pp. 495-512. In, Methodology of plant eco-physiology, F.E. Eckardt, ed. UNESCO, Arid Zone Research 25. 531p.
- Baumgartner, A. 1967. Energetic bases for differential vaporization from forest and agricultural lands, pp. 381-389. In, Forest Hydrology, W.E. Sopper and H.W. Lull, eds. Pergamon Press, New York. 813p.
- Beal, W.J. 1904. Some of the changes now taking place in a forest of oak openings. Michigan Acad. Sci. Ann. Rep. 4: 107-108.
- Beard, J.S. 1953. The savanna vegetation of northern tropical America. Ecol. Monogr. 23: 149-215.
- Beers, T.W. and C.I. Miller. 1964. Point sampling: Research results. Theory and applications. Purdue Univ. Agr. Exp. Sta. Res. Bull. 786. 56p.
- Bergman, H.F. 1924 (1923). The composition of climax plant formations in Minnesota. Michigan Acad. Sci. Arts and Letters, Papers 3: 51-60.
- Bergman, H.F. and H. Stallard. 1916. The development of climax formations in northern Minnesota. Minnesota Bot. Stud. 4: 333-378.
- Berkner, L.V. and L.C. Marshall. 1965. History of major atmospheric components. Proc. Nat. Acad. Sci. 53(6): 1215-1226.
- Billings, W.D. 1952. The environmental complex in relation to plant growth and distribution. Quart. Rev. Biol. 27: 251-265.
- Billings, W.D. and R.J. Morris. 1951. Reflection of visible and infrared radiation from leaves of different ecological groups. Amer. J. Bot. 38: 327-331.
- Bird, R.D. 1930. Biotic communities of the aspen parkland of central Canada. Ecology 11: 356-442.
- Birkebak, R. and R. Birkebak. 1964. Solar radiation characteristics of tree leaves. Ecology 45: 646-649.

- Blake, G.R., E.R. Allred, C.H.M. van Bavel, and F.D. Whisler. 1960. Agricultural drought and moisture excesses in Minnesota. Univ. Minnesota Agr. Exp. Sta. Tech. Bull. 235. 36p.
- Bolin, B. 1970. The carbon cycle. Sci. Amer. 223(3): 124-132.
- Borchert, J.R. 1950. The climate of the central North American Grassland. Ass. Amer. Geogr., Ann. 40: 1-39.
- Borchert, J.R. 1971. The dust bowl in the 1970's. Ass. Amer. Geogr., Ann. 61: 1-22.
- Bourdo, E.A. 1956. A review of the General Land Office Survey and of its use in quantitative studies of former forests. Ecology 37: 754-768.
- Bourne, A. 1820. On the prairies and barrens of the west. Amer. J. Sci., Arts 2: 30-34.
- Bowen, I.S. 1926. The ratio of heat losses by conduction and by evaporation from any water surface. Phys. Rev. 27: 779-787.
- Bowman, I. 1914. Forest physiography. Physiography of the United States and principles of soils in relation to forestry. John Wiley and Sons, Inc., New York 759p.
- Braun, E.L. 1938. Deciduous forest climaxes. Ecology 19: 515-522.
- Braun, E.L. 1947. Development of the deciduous forests of eastern North America. Ecol. Monogr. 17: 211-219.
- Braun, E.L. 1950 (1967). Deciduous forests of eastern North America. Hafner Publ. Co., New York, 596p.
- Braun, E.L. 1951. Plant distribution in relation to the glacial boundary. Ohio J. Sci. 51: 139-146.
- Bray, E.C. 1962. A million years in Minnesota. The glacial story of the state. The Sci. Mus., St. Paul. 49p.
- Bray, J.R. 1955. The savanna vegetation of Wisconsin and an application of the concepts order and complexity to the field of ecology. Ph.D. Thesis, Univ. Wisconsin. 174p., plus appendix.
- Bray, J.R. 1968. Glaciation and solar activity since the fifth century BC and the solar cycle. Nature 220: 672-674.

- Bray, J.R. 1971. Vegetation distribution, tree growth and crop success in relation to recent climatic change. *Adv. Ecol. Res.* 7: 177-233.
- Bray, J.R., J.E. Sanger, and A.L. Archer. 1966. The visible albedo of surfaces in central Minnesota. *Ecology* 47: 524-531.
- Bray, R.H. and L.T. Kurtz. 1945. Determination of total, organic, and available forms of phosphorus in soils. *Soil Sci.* 59: 39-45.
- Bremner, J.M. 1965. Total nitrogen, pp. 1149-1178. In, *Methods of soil analysis, Part 2. Chemical and microbiological properties.* C.A. Black, ed. Amer. Soc. Agron., Madison, Wisconsin. (No. 9 in the series *Agronomy*). 1572p., plus indices.
- Brinkman, R.T. 1969. Dissociation of water vapor and evolution of oxygen in the terrestrial atmosphere. *J. Geophys. Res.* 74: 5355-5368.
- Brooks, C.E.P. 1949. *Climate through the ages. A study of the climatic factors and their variations.* McGraw-Hill Book Co., New York. 395p.
- Brooks, F.A. 1961. Need for measuring horizontal gradients in determining vertical eddy transfers of heat and moisture. *J. Meteorol.* 18: 589-596.
- Bryson, R.A. and W.M. Wendland. 1967. Tentative climatic patterns for some late glacial and post-glacial episodes in central North America, pp. 271-298. In, *Life, land and water*, W.J. Mayer-Oakes, ed. (Proc. 1966 Conf. Environ. Stud. Glacial Lake Agassiz Region. Univ. Manitoba Dept. Anthropol. Occ. Paper No. 1.) Univ. Manitoba Press, Winnipeg. 416p.
- Buckman, H.O. and N.C. Brady. 1962. *The nature and property of soils.* (6th ed.) Macmillan Co., New York. 567p.
- Budyko, M.I. 1948 (1963). *Evaporation under natural conditions.* GIMIZ. Gidrometeorologicheskoe Izdalel'stvo, Leningrad. (Transl. OTS 63-11061.) 130p.
- Budyko, M.I. 1956. *The heat balance of the earth's surface.* Gidrometeorologicheskoe Izdalel'stvo, Leningrad. 255p. (Transl. U.S. Dept. Commerce, Office of Climatology, 1958. 259p.)

- Buell, M.F. and H.F. Buell. 1959. Aspen invasion of prairie. Bull. Torrey Bot. Club 86: 264-265.
- Buell, M.F. and J.E. Cantlon. 1951. A study of two forest stands in Minnesota with an interpretation of the prairie-forest margin. Ecology 32: 294-316.
- Buell, M.F. and V. Facey. 1960. Forest-prairie transition west of Itasca Park, Minnesota. Bull. Torrey Bot. Club 87: 46-58.
- Bunge, M. 1967. Scientific research I. The search for system. Springer-Verlag, New York. 536p.
- Cain, S.A. 1944. Foundations of plant geography. Harper and Brothers, New York. 556p.
- Campbell, J.T. 1886a. Causes of forest rotation. Amer. Natur. 20: 521-527.
- Campbell, J.T. 1886b. Causes of forest rotation. Amer. Natur. 20: 851-856.
- Carpenter, J.R. 1940. The grassland biome. Ecol. Monogr. 10: 617-684.
- Carpenter, W.M. 1839. Miscellaneous notices in Opelousas, Attakapas, &c. Amer. J. Sci. Arts 35: 344-346.
- Catlin, G. 1850. Illustrations of the manners, customs, and condition of the North American Indians with letters and notes written during eight years of travel and adventure among the wildest and most remarkable tribes now existing. Henry G. Bohn, London. 9th ed. Two volumes. Vol. 1: 264p. Vol. 2: 266p.
- Caton, J.D. 1876. Origin of the prairies. Fergus' Histor. Ser. 1(7): 31-55.
- Chaney, R.W. and M.K. Elias. 1938 (1936). Late Tertiary floras from the high plains, pp. 1-46. In, Miocene and Pliocene floras of western North America. Carnegie Inst. Washington, Washington, D.C. Pub. No. 476. 272p.
- Chang, J. 1968. Climate and agriculture. An ecological survey. Aldine Publ. Co., Chicago. 304p.
- Chapman, H.D. and P.F. Pratt. 1961. Methods of analysis for soils, plants and waters. Div. Agr. Sci., Univ. California Riverside. 309p.

Chavannes, E.A. 1940. The steep prairies of southern Wisconsin and their invasion by forest. Ph.D. Thesis, Univ. Wisconsin. 56p., plus intro. and bibliog.

Chavannes, E. 1941. Written records of forest succession. Sci. Monthly 53: 76-80.

Christy, M. 1892. Why are the prairies treeless? Proc. Roy. Geogr. Soc. and Monthly Rec. Geogr. 14: 78-100.

Clark, A.H. 1956. The impact of exotic invasion on the remaining New World mid-latitude grasslands, pp. 737-762. In, Man's role in changing the face of the earth, W.L. Thomas, Jr., ed. Univ. Chicago Press, Chicago. 1193p.

Clarke, G.L. 1954. Elements of ecology. John Wiley and Sons, Inc., New York. 534p.

Clements, F.E. 1916. Plant succession. An analysis of the development of vegetation. Carnegie Inst. Washington Pub. No. 242. Washington, D.C. 512p.

Clements, F.E. and R.W. Chaney. 1936. Environment and life in the Great Plains. Carnegie Inst. Washington Suppl. Pubs. 24. 54p.

Clements, F.E., C.O. Rosendahl, and F.K. Butters. 1912. Minnesota trees and shrubs. An illustrated manual of the native and cultivated woody plants of the state. Minnesota Geol. and Natur. Hist. Surv., Rep. Bot. Surv., 9. 314p.

Cloud, P.E., Jr. 1968. Atmospheric and hydrospheric evolution on the primitive earth. Science 160: 729-736.

Cloud, P. and A. Gibor. 1970. The oxygen cycle. Sci. Amer. 223(3): 110-123.

Coats, R.N. 1967. Evaluation of an ecological nutrient gradient in the central pine section of Minnesota. M.S. Thesis, Univ. Minnesota, Minneapolis. 120p.

Coats, R.N., W.A. Geyer, and E.I. Sucoff. 1968. Synecological light coordinates: a verification by light measurements. Minnesota Forest. Res. Notes No. 199. 4p.

Conard, H.S. 1951. The background of plant ecology. (Transl. The plant life of the Danube Basin, Anton Kerner, 1863.) Iowa State Coll. Press, Ames. 238p.

Conway, V.M. 1948. Von Post's work on climatic rhythms. *New Phytol.* 47: 220-237.

Cooper, J.F. 1848 (1900). *Oak openings*. R.F. Fenno and Co., New York. 412p.

Cooper, J.G. 1859. On the distribution of the forests and trees of North America, with notes on its physical geography, pp. 246-280. *In*, Smithsonian Inst. Ann. Rep. for 1858. Washington, D.C.

Costello, D.F. 1969. *The prairie world*. Thomas Y. Crowell Co., New York. 242p.

Cottam, G. 1949. The phytosociology of an oak woods in southwestern Wisconsin. *Ecology* 30: 271-287.

Coulombe, H.N. and J. Brown. 1970. The synthesis and modeling of the Barrow, Alaska, ecosystem. *Int. Union Conserv. Nature and Natur. Res.*, New Ser. 16: 44-49.

Curtis, J.T. 1950. *Plant ecology work book; laboratory, field, and reference manual*. Burgess Pub. Co., Minneapolis. 39(64)p.

Curtis, J.T. 1956. The modification of mid-latitude grasslands and forests by man, pp. 721-736. *In*, *Man's role in changing the face of the earth*, W.L. Thomas, Jr., ed. Univ. Chicago Press, Chicago. 1193p.

Curtis, J.T. 1959. *The vegetation of Wisconsin*. Univ. Wisconsin Press, Madison. 657p.

Curtis, J.T. and H.C. Greene. 1949. A study of relic Wisconsin prairies by the species-presence method. *Ecology* 30: 83-92.

Curtis, J.T. and R.P. McIntosh. 1951. An upland continuum in the prairie-forest border region of Wisconsin. *Ecology* 32: 476-496.

Curtis, J.T. and M.L. Partch. 1948. Effect of fire on the competition between blue grass and certain prairie plants. *Amer. Midland Natur.* 39: 437-443.

Cushing, E.J. 1965. Problems in the Quaternary phytogeography of the Great Lakes region, pp. 403-416. *In*, *Quaternary of the United States*, H.E. Wright, Jr. and D.G. Frey, eds. Princeton Univ. Press, Princeton. 922p.

- Cushing, E.J. 1967. Late-Wisconsin pollen stratigraphy and the glacial sequence in Minnesota, pp. 59-88. In, Quaternary paleoecology, E.J. Cushing and H.E. Wright, Jr., eds. (Vol. 7 of the Proc. VII Congr. Int. Ass. Quaternary Res.) Yale Univ. Press, New Haven. 433p.
- Dana, J.D. 1865. On the origin of prairies. Amer. J. Sci. Arts, Second Ser. 40: 293-304.
- Daubenmire, R.F. 1936. The "Big Woods" of Minnesota: its structure, and relation to climate, fire, and soils. Ecol. Monogr. 6: 233-268.
- Daubenmire, R. 1968a. Ecology of fire in grasslands. Adv. Ecol. Res. 5: 209-266.
- Daubenmire, R. 1968b. Plant communities. A textbook of plant synecology. Harper and Row, Publ., New York. 300p.
- Dawson, G.M. 1875. Report on the geology and resources of the region in the vicinity of the forty-ninth parallel, from the Lake of the Woods to the Rocky Mountains, with lists of plants and animals collected, and notes on the fossils. Rep. to Brit. North Amer. Boundary Comm. Dawson Brothers, Montreal. 387p.
- Day, G.M. 1953. The Indian as an ecological factor in the northeastern forest. Ecology 34: 329-346.
- Day, P.R. 1965. Particle fractionation and particle size analysis, pp. 545-567. In, Methods of soil analysis, Part 1, C.A. Black, ed. Amer. Soc. Agron., Madison, Wisconsin. (No. 9 in the series Agronomy). 1572p., plus indices.
- Deacon, E.L. 1950. The measurement and recording of the heat flux into the soil. Quart. J. Roy. Meteorol. Soc. 76: 479-483.
- Deacon, E.L. and W.C. Swinbank. 1958. Comparison between momentum and water vapor transport, pp. 38-41. In, Climatology and micrometeorology. (Proc. Canberra Symp.). UNESCO. Arid Zone Research 11. 355p.
- Deevey, E.S., Jr. 1949. Pleistocene research. 3. Biogeography of the Pleistocene. Part 1: Europe and North America. Bull. Geol. Soc. Amer. 60: 1315-1416.

- Denmead, O.T. 1964. Evaporation sources and apparent diffusivities in a forest canopy. *J. Appl. Meteorol.* 3: 383-389.
- Denmead, O.T. 1968. The energy balance of plant communities, pp. 71-105. *In*, *Agricultural Meteorology*, Vol. 1. (Proc. WMO Seminar, Melbourne, Australia, 1966). Bureau Meteorology, Australia. 263p.
- Denmead, O.T. 1970. Transfer processes between vegetation and air: Measurement, interpretation and modelling, pp. 149-164. *In*, *Prediction and measurement of photosynthetic productivity*. (Proc. IBP/PP Tech. Meeting, Sept., 1969, I. Málek, chm.). Centre for Agr. Pub. and Doc., Wageningen. 632p.
- Deters, M.E. and H. Schmitz. 1936. Drouth damage to prairie shelterbelts in Minnesota. *Minnesota Agr. Exp. Sta. Bull.* 329. 28p.
- DeVries, D.A. 1963. Thermal properties of soils, pp. 210-235. *In*, *Physics of the plant environment*, W.R. Van Wijk, ed. John Wiley and Sons, Inc., New York. 382p.
- Dice, L.R. 1952. *Natural communities*. Univ. Michigan Press, Ann Arbor. 547p.
- Dix, R.L. and J.E. Butler. 1954. The effects of fire on a dry, thin-soil prairie in Wisconsin. *J. Range Manage.* 7: 265-268.
- Dix, R.L. and F.E. Smeins. 1967. The prairie, meadow, and marsh vegetation of Nelson County, North Dakota. *Can. J. Bot.* 45: 21-58.
- Dixon, W.J., ed. 1971. BMD. Biomedical computer programs. Univ. California Pub. Automatic Computation No. 2. Univ. California Press, Berkeley. 600p.
- Doing, H. 1969. Sociological species groups. *Acta Bot. Neerl.* 18: 398-400.
- Dorf, E. 1960. Climatic changes of the past and present. *Amer. Sci.* 48: 341-364.
- Douglass, A.E. 1919. Climatic cycles and tree-growth. [Vol. 1]. A study of the annual rings of trees in relation to climate and solar activity. Carnegie Inst. Washington, Washington, D.C. Pub. No. 289. 127p.

- Douglass, A.E. 1928. Climatic cycles and tree-growth.
Vol. 2. A study of the annual rings of trees in relation to climate and solar activity. Carnegie Inst. Washington, Washington, D.C. Pub. No. 289. 166p.
- Douglass, A.E. 1929. The secret of the southwest solved by talkative tree rings. Nat. Geogr. Mag. 56: 736-770.
- Douglass, A.E. 1936. Climatic cycles and tree growth.
Vol. 3. A study of cycles. Carnegie Inst. Washington, Washington, D.C. Pub. No. 289. 171p.
- Douglass, J.E. 1967. Effects of species and arrangement of forests on evapotranspiration, pp. 451-461. In, Forest hydrology, W.E. Sopper and H.W. Lull, eds. Pergamon Press, New York. 813p.
- Drew, L.A. 1973. Bulk density estimation based on organic matter content of some Minnesota soils. Minnesota Forest. Res. Notes No. 243. 4p.
- Drummond, A.J. 1964. Comments on "sky radiation measurement and correction". J. Appl. Meteorol. 3: 810-811.
- Duvigneaud, P. and S. Denaeyer-DeSmet. 1970. Phytogéochimie des groupes écosociologiques forestiers de Haute-Belgique. 1. Essai de classification phytochimique des espèces herbacées. Oecologia Plantarum 5: 1-31.
- Dyer, A.J. 1961. Measurements of evaporation and heat transfer in the lower atmosphere by an automatic eddy-correlation technique. Quart. J. Roy. Meteorol. Soc. 87: 401-412.
- Dyer, A.J. 1963. The adjustment of profiles and eddy fluxes. Quart. J. Roy. Meteorol. Soc. 89: 276-280.
- Dyer, A.J. 1967. The turbulent transport of heat and water vapour in an unstable atmosphere. Quart. J. Roy. Meteorol. Soc. 93: 501-508.
- Dyksterhuis, E.J. 1957. The savanna concept and its use. Ecology 38: 435-442.
- Eiseley, L.C. 1954. Man the fire-maker. Sci. Amer. 191(3): 52-57.
- Elias, M.K. 1935. Tertiary grasses and other prairie vegetation from High Plains of North America. Amer. J. Sci., Ser. 5, 29: 24-33.

- Elson, J.A. 1967. Geology of Glacial Lake Agassiz, pp. 37-95. In, Life, land and water, W.J. Mayer-Oakes, ed. (Proc. 1966 Conf. Environ. Stud. Glacial Lake Agassiz Region. Univ. Manitoba Dept. Anthropol. Occ. Paper No. 1). Univ. Manitoba Press, Winnipeg. 416p.
- Elwell, H.M., H.A. Daniel, and F.A. Fenton. 1941. The effects of burning pasture and woodlands vegetation. Oklahoma Agr. Exp. Sta. Bull. B-247. 14p.
- Engelmann, H. 1863. Remarks upon the causes producing the different characters of vegetation known as Prairies, Flats, and Barrens in southern Illinois, with special reference to observations made in Perry and Jackson counties. Amer. J. Sci. Arts, Second Ser., 36: 384-396.
- Evans, R.T., ed. 1963. Environmental control of plant growth. Academic Press, New York. 449p.
- Ewing, J. 1924. Plant successions of the brush-prairie in northwestern Minnesota. J. Ecol. 12: 238-266.
- Federer, C.A. 1970. Measuring forest evapotranspiration--theory and problems. U.S.D.A., Forest Service. Northeastern Forest Exp. Sta., Res. Paper NE - 165. 25p.
- Fendler, A. 1866. On prairies. Amer. J. Sci. Arts, Ser. II, 41: 154-158.
- Fenneman, N.M. 1938. Physiography of eastern United States. McGraw-Hill Book Co., Inc., New York. 714p.
- Feyerham, A.M., L.D. Bark, and W.C. Burrows. 1966. Probabilities of sequences of wet and dry days in Minnesota. Kansas State Univ. Agr. Exp. Sta. Tech. Bull. 139j. (North Central Regional Res. Pub. 161). 55p.
- Finn, R.F. 1953. Mineral content of leaves clue to white oak site quality. U.S. Dept. Agr. Central States Forest Exp. Sta. Tech. Pap. 135. 12p.
- Fitzpatrick, E.A. and W.R. Stern. 1973. Net radiation estimated from global solar radiation, pp. 403-410. In, Plant response to climatic factors, R.O. Slayter, ed. UNESCO, Ecology and Conservation, 5. (Proc. Uppsala Symp., 1970). 574p.
- Flaccus, E. 1965. Distribution in Minnesota of Acer saccharum, Tilia americana, and Betula lutea. Minnesota Acad. Sci. Proc. 32: 95-97.

- Flint, R.F. 1947. Glacial geology and the Pleistocene Epoch. John Wiley and Sons, Inc., New York. 589p.
- Flint, R.F. 1953. Evidence from glacial geology as to climatic variations, pp. 165-178. In, Climatic changes. Evidence, causes, and effects, H. Shapley, ed. Harvard Univ. Press, Cambridge. 318p.
- Flint, R.F. 1971. Glacial and Quaternary geology. John Wiley and Sons, Inc., New York. 892p.
- Flint, R.F. and F. Brandtner. 1961. Outline of climatic fluctuation since the last interglacial age. New York Acad. Sci., Ann. 95: 457-460.
- Flohn, H. 1961. Man's activity as a factor in climatic change. New York Acad. Sci., Ann. 95: 271-281.
- Flohn, H. 1969. Climate and weather. McGraw-Hill Book Co., New York. 253p.
- ~~Foster~~, J.W. 1869. The Mississippi valley: its physical geography, including sketches of the topography, botany, climate, geology, and mineral resources; and of the progress of development in population and material wealth. S.C. Griggs and Co., Chicago. 443p.
- Fowells, H.A. 1965. Silvics of forest trees of the United States. U.S. Dept. Agr., Forest Serv., Washington, D.C. 762p.
- Fowler, W.C. 1965. The energy budget and its use in estimating evapotranspiration. Proc. Soc. Amer. Forest. 1964: 101-104.
- Fritschen, L.J. 1963. Construction and evaluation of a miniature net radiometer. J. Appl. Meteorol. 2: 165-172.
- Fritschen, L.J. 1965a. Accuracy of evapotranspiration determinations by the Bowen ratio method. Bull. Inter. Ass. Sci. Hydrol. 10(2): 38-48.
- Fritschen, L.J. 1965b. Miniature net radiometer improvements. J. Appl. Meteorol. 4: 528-532.
- Fritschen, L.J. 1966. Evapotranspiration rates of field crops determined by the Bowen ratio method. Agron. J. 58: 339-342.

- Fuchs, M. and C.B. Tanner. 1968. Calibration and field test of soil heat flux plates. *Soil Sci. Soc. Amer. Proc.* 32: 326-328.
- Fuller, G.D. 1914. Evaporation and soil moisture in relation to the succession of plant associations. *Bot. Gaz.* 58: 193-234.
- Fuller, G.D. 1923. An edaphic limit to forests in the prairie region of Illinois. *Ecology* 4: 135-140.
- Funk, J.P. 1959. Improved polythene-shielded net radiometer. *J. Sci. Instruments* 36: 267-270.
- Gagnon, J.D. 1964. Relationship between site index and foliage nitrogen at two crown levels for mature black spruce. *Forest. Chron.* 40: 169-174.
- Gagnon, J.D., A. Lafond, and L.P. Amiot. 1958. Mineral nutrient content of some forest plant leaves and of the humus layer as related to site quality. *Canad. J. Bot.* 36: 209-220.
- Garretson, M.S. 1938. The American bison. New York Zool. Soc., New York. 254p.
- Gaskill, A. 1906. Why prairies are treeless. *Proc. Soc. Amer. Forest.* 1(3): 158-178.
- Gates, D. 1962. Energy exchange in the biosphere. Harper and Row, Publ., New York. 151p.
- Gates, D.M. 1965. Characteristics of soil and vegetated surfaces to reflected and emitted radiation, pp. 573-600. *Symposium on Remote Sensing of Environment*, Univ. Michigan, Inst. Sci. Tech., Proc. 3. 821p.
- Gates, D.M. 1968. Toward understanding ecosystems. *Adv. Ecol. Res.* 5: 1-35.
- Gates, D.M. and R.J. Hanks. 1967. Plant factors affecting evapotranspiration, pp. 506-521. *In*, *Irrigation of agricultural lands*, R.M. Hagen, H.R. Haise, and T.W. Edminster, eds. *Amer. Soc. Agron., Madison, Wisconsin. Agronomy II.* 1180p.
- Gates, D.M., H.J. Keegan, J.C. Schleter, and V.R. Weidner. 1965. Spectral properties of plants. *Appl. Optics* 4: 11-20.

- Geiger, R. 1965 (1957). The climate near the ground. Harvard Univ. Press, Cambridge, Massachusetts. 494p.
- Gerloff, G.C., D.G. Moore, and J.T. Curtis. 1964. Mineral content of native plants of Wisconsin. Univ. Wisconsin Agr. Exp. Sta. Res. Rep. 14. 27p.
- Gerloff, G.C., D.G. Moore, and J.T. Curtis. 1966. Selective absorption of mineral elements by native plants of Wisconsin. Plant and Soil 25: 393-405.
- Gleason, H.A. 1909. Some unsolved problems of the prairies. Bull. Torrey Bot. Club 36: 265-271.
- Gleason, H.A. 1912. An isolated prairie grove and its phytogeographical significance. Bot. Gaz. 53: 38-49.
- Gleason, H.A. 1913. The relation of forest distribution and prairie fires in the Middle West. Torrey 13: 173-181.
- Gleason, H.A. 1922. The vegetational history of the Middle West. Ass. Amer. Geogr., Ann. 12: 39-85.
- Gleason, H.A. 1926. The individualistic concept of the plant association. Bull. Torrey Bot. Club 53: 7-26.
- Gleason, H.A. 1939. The individualistic concept of the plant association. Amer. Midland Natur. 21: 92-110.
- Gleason, H.A. and A. Cronquist. 1963. Manual of vascular plants of Northeastern United States and adjacent Canada. D. VanNostrand Co., Inc., Princeton, New Jersey. 810p.
- Gleason, H.A. and A. Cronquist. 1964. The natural geography of plants. Columbia Univ. Press, New York. 420p.
- Good, R.E. and N.F. Good. 1971. Vegetation of a Minnesota prairie and a comparison of methods. Amer. Midland Natur. 85: 228-231.
- Good, R. 1964 (1947). The geography of the flowering plants. John Wiley and Sons, Inc., New York. 518p.
- Goodall, D.W. and F.G. Gregory. 1947. Chemical composition of plants as an index of their nutritional status. Imp. Bur. Hort. and Plantation Crops, Tech. Comm. No. 17. 167p.

- Gordon, A.G. 1964. The nutrition and growth of ash, Fraxinus excelsior, in natural stands in the English lake district as related to edaphic site factors. J. Ecol. 52: 169-187.
- Gould, F.W. 1941. Plant indicators of original Wisconsin prairies. Ecology 22: 427-429.
- Gove, P.B., ed. 1967. Webster's third new international dictionary of the English language. Unabridged. G.&C. Merriam Co., Springfield, Massachusetts. 2662p.
- Gray, A. 1857. Statistics of the flora of the northern United States. Amer. J. Sci. Arts, Second Ser., 23: 369-403.
- Gray, A. 1885. Characteristics of the North American flora. Nature 31: 232-235; 253-255.
- Green, P.A. 1950. Ecological composition of high prairie relics in Rock County, Wisconsin. Wisconsin Acad. Sci. Arts Letters, Trans. 40(1): 159-172.
- Grigal, D.F. 1968. The relationship between plant communities and soils in northeastern Minnesota. Ph.D. Thesis, Univ. Minnesota, Minneapolis. 288p.
- Griggs, R.F. 1914. Observations on the behavior of some species on the edges of their ranges. Bull. Torrey Bot. Club 41: 25-49.
- Grosenbaugh, L.R. 1952. Plotless timber estimates--new, fast, easy. J. Forest. 50: 32-37.
- Gunderson, H.L. and J.R. Beer. 1953. The mammals of Minnesota. Minnesota Univ. Mus. Natur. Hist. Occ. Paper No. 6. 190p.
- Habeck, J.R. 1959. A phytosociological study of the upland forest communities in the central Wisconsin sand plain area. Wisconsin Acad. Sci. Arts Letters, Trans. 48: 31-48.
- Hadley, E.B. 1970. Net productivity and burning responses of native eastern North Dakota prairie communities. Amer. Midland Natur. 84: 121-135.
- Hall, C.W. 1903. Geography and geology of Minnesota. Vol. 1. Geography of Minnesota. H.W. Wilson Co., Minneapolis. 299p.
- Hall, N., V.S. Russell, and C.D. Hamilton. 1958. Trees and microclimate, pp. 259-264. In, Climatology and micro-climatology. (Proc. Canberra Symposium). UNESCO. Arid Zone Research 11. 355p.

- Haines, F. 1970. The buffalo. T.Y. Crowell Co., New York. 242p.
- Hanson, H.C. 1922. Prairie inclusions in the deciduous forest climax. Amer. J. Bot. 9: 330-337.
- Hanson, H.C. 1962. Dictionary of ecology. Phil. Library, Inc., New York. 382p.
- Hanson, L.D. 1973. General characteristics of Minnesota soils, pp. 130-138. In, Soils, soil management and fertilizer monographs. Univ. Minnesota Agr. Ext. Serv. Spec. Rep. 24. 157p.
- Harshberger, J.W. 1911. Phytogeographic survey of North America. Wilhelm Engelmann, Leipzig. 790p.
- Harvey, E.M. 1913. Evaporation and soil moisture on the prairies of Illinois. Illinois Acad. Sci., Trans. 6: 92-99.
- Harvey, L.H. 1908a. Floral succession in the prairie-grass formation of southeastern South Dakota. Serotinal and autumnal aspects. Bot. Gaz. 46: 277-298.
- Harvey, L.H. 1908b. Floral succession in the prairie-grass formation of southeastern South Dakota. The prevernal, vernal, and estival aspects. Bot. Gaz. 46: 81-108.
- Hatfield, H.S. and F.J. Wilkins. 1950. New heat-flow meter. J. Sci. Instruments, London 27: 1-3.
- Hay, O.P. 1878. An examination of Prof. Leo Lesquereux's theory of the origin and formation of prairies. Amer. Natur. 12: 299-305.
- Henry, D.G. 1972. Foliar nutrient patterns in the subordinate vegetation of six Minnesota forest stands. M.S. Thesis, Univ. Minnesota, Minneapolis. 222p.
- Hensel, R.L. 1923. Effect of burning on vegetation in Kansas pastures. J. Agr. Res. 23: 631-643.
- Holch, A.E. 1931. Development of roots and shoots of certain deciduous tree seedlings in different forest soils. Ecology 12: 259-298.
- Horowitz, J.L. 1969. An easily constructed shadow-band for separating direct and diffuse solar radiation. Solar Energy 12: 543-545.

- Horton, J.S. 1973. Evapotranspiration and water research as related to riparian and phreatophyte management. An abstract bibliography. U.S.D.A., Forest Service, Misc. Pub. No. 1234. 192p.
- Hoyle, M.C. and D.L. Mader. 1964. Relationships of foliar nutrients to growth of red pine in western Massachusetts. *Forest Sci.* 10: 337-347.
- Hulbert, L.C. 1969. Fire and litter effects in undisturbed bluestem prairie in Kansas. *Ecology* 50: 874-877.
- Jackson, M.L. 1958. Soil chemical analysis. Prentice-Hall, Inc., Englewood Cliffs, New Jersey. 498p.
- Jelgersma, S. 1962. A late-glacial pollen diagram from Madelia, south-central Minnesota. *Amer. J. Sci.* 260: 522-529.
- Jenny, H. 1941. Factors of soil formation. A system of quantitative pedology. McGraw-Hill Book Co., Inc., New York. 281p.
- Johnsen, S.J., W. Dansgaard, and H.B. Clausen. 1970. Climatic oscillations 1200-2000 A.D. *Nature* 227: 482-483.
- Jones, G. 1838. Some observations in Holland, connected with our prairie region. *Dry prairies. Amer. J. Sci. Arts* 33: 226-230.
- Just, T. 1959. Postglacial vegetation of the north-central United States: a review. *J. Geol.* 67: 228-238.
- Keeling, C.D. 1970. Is carbon dioxide from fossil fuel changing man's environment? *Amer. Phil. Soc., Proc.* 114(1): 10-17.
- Kersten, M.S. 1949. Thermal properties of soils. *Univ. Minnesota, Inst. Tech., Eng. Exp. Sta. Bull.* 28. 227p.
- King, P.B. 1965. Tectonics of Quaternary time in Middle North America, pp. 831-870. In, *The Quaternary of the United States*, H.E. Wright, Jr. and D.G. Frey, eds. Princeton Univ. Press, Princeton. 922p.
- Kirsch, L.M. and A.D. Kruse. 1973. Prairie fires and wildlife. *Proc. Ann. Tall Timbers Fire Ecol. Conf.* 12: 289-303.
- Knoerr, K.R. 1965. Partitioning of the radiant heat load by forest stands. *Proc. Soc. Amer. Forest.* 1964: 105-109.

- Knoerr, K.R. 1967. Contrasts in energy balances between individual leaves and vegetated surfaces, 391-401. In, Forest hydrology, W.E. Sopper and H.W. Lull, eds. Pergamon Press, New York. 813p.
- Kohnke, H. and G.A. Nakshabandi. 1964. Heat transfer in soils. Trans. 8th Intern. Congr. Soil Sci. Bucharest. 2: 185-193.
- Komarek, E.V., Sr. 1964. The natural history of lightning. Proc. Ann. Tall Timbers Fire Ecol. Conf. 3: 139-183.
- Kraus, E.B. 1961. Physical aspects of deduced and actual climatic change. New York Acad. Sci., Ann. 95: 225-234.
- Krieth, F. 1965 (1967). Principles of heat transfer. Second ed. International Textbook Co., Scranton, Pennsylvania. 620p.
- Kucera, C.L. and J.H. Ehrenreich. 1962. Some effects of annual burning on central Missouri prairie. Ecology 43: 334-336.
- Kucera, C.L. and M. Koelling. 1964. The influence of fire on composition of central Missouri prairie. Amer. Midland Natur. 72: 142-147.
- Kucera, C.L. and R.E. McDermott. 1955. Sugar maple-basswood studies in the forest-prairie transition of central Missouri. Amer. Midland Natur. 54: 495-503.
- Küchler, A.W. 1964. Potential natural vegetation of the conterminous United States. Amer. Geogr. Soc., Spec. Pub. No. 36. 155p., plus map.
- Kuehnast, E.L. 1972 (1959). Climate of Minnesota. Climatology of the United States No. 60-21. U.S. Dept. Com., Nat. Oceanic and Atmos. Admin. 20p.
- Kuehnast, E.L. 1973. Personal communication.
- Kulp, J.L. 1953. Climatic changes and radioisotope dating, pp. 201-208. In. Climatic change. Evidence, causes, and effects, H. Shapley, ed. Harvard Univ. Press, Cambridge. 318p.
- Kupsch, W.O. 1967. Postglacial uplift -- A review, pp. 155-186. In, Life, land and water, W.J. Mayer-Oakes, ed. (Proc. 1966 Conf. Environ. Stud. Glacial Lake Agassiz Region. Univ. Manitoba Dept. Anthropol. Occ. Paper No. 1). Univ. Manitoba Press, Winnipeg. 416p.

- Kurmis, V. 1969. Dynamics of advance reproduction in upland forest communities in Itasca State Park, Minnesota. Ph.D. Thesis, Univ. Minnesota, Minneapolis. 166p.
- Lamb, H.H. 1969. The new look of climatology. *Nature* 223: 1209-1215.
- Landsberg, H.E. 1970. Man-made climatic changes. *Science* 170: 1265-1274.
- Lane, G.H. 1931. A preliminary pollen analysis of the east McCulloch peat bed. *Ohio J. Sci.* 31: 165-171.
- Lawrence, D.B. 1950. Glacier fluctuation for six centuries in southeastern Alaska and its relation to solar activity. *Geogr. Rev.* 40: 191-223.
- Lawrence D.B. 1958. Glaciers and vegetation in southeastern Alaska. *Amer. Sci.* 46: 89-122.
- Lawrence, D.B. 1973. Personal communication of data from the National Oceanic and Atmospheric Administration, World Data Center A, Upper Atmosphere Geophysics, Boulder, Colorado. Wolf Mean Monthly Sunspot Numbers, 1947-1970.
- Lawrence, D.B., R.E. Schoenike, A. Quispel, and G. Bond. 1967. The role of Dryas drummondi in vegetation development following ice recession at Glacier Bay, Alaska, with special reference to its nitrogen fixation by root nodules. *J. Ecol.* 55: 793-813.
- Lee, S.C. 1924. Factors controlling forest succession at Lake Itasca, Minnesota. *Bot. Gaz.* 78: 129-174.
- Leighton, M.M. 1960. The classification of the Wisconsin glacial stage of north central United States. *J. Geol.* 68: 529-552.
- Lemon, E. 1963. Energy and water balance of plant communities, pp. 55-78. In, Environmental control of plant growth, L.T. Evans, ed. Academic Press, New York. 449p.
- Lemon, E. 1965. Micrometeorology and physiology of plants in their natural environment, pp. 203-227. In, Plant physiology. A treatise. Vol. IV A: Metabolism: Organic nutrition and nitrogen metabolism, F.C. Steward, ed. Academic Press, New York. 731p.
- Lesquereux, L. 1865a. On the origin and formation of prairies. *Amer. J. Sci. Arts, Second Ser.*, 39: 317-327.

- Lesquereux, L. 1865b. On the origin and formation of prairies. Amer. J. Sci. Arts, Second Ser., 40: 23-31.
- Lettau, H.H. 1957. Computation of heat budget constituents of the Earth/Air interface, pp. 305-327. In, Exploring the atmosphere's first mile, Vol. 1, H.H. Lettau and B. Davidson, eds. Instrumentation and data evaluation. (Proc. Great Plains Turbulence Field Program, 1953). (U.S. Air Force Cambridge Research Center, Geophysics Research Directorate, Air Research and Development Command). Pergamon Press, New York. 376p.
- Leverett, F. and F.W. Sardeson. 1932. Quaternary geology of Minnesota and parts of adjacent states. U.S. Geol. Surv. Prof. Pap. 161. 149p.
- Leyton, L. 1956. The relation between growth and mineral composition of foliage of Japanese larch (Larix leptolepis, Murr.). Plant and Soil 7: 167-177.
- Leyton L. 1958. The relationship between the growth and mineral nutrition of conifers, pp. 323-345. In, The physiology of forest trees, K.V. Thimann, ed. Ronald Press, New York. 678p.
- Leyton, L. and K.A. Armson. 1955. Mineral composition of the foliage in relation to the growth of Scots pine. Forest Sci. 1: 210-218.
- Lieth, H. 1963. The role of vegetation in the carbon dioxide content of the atmosphere. J. Geophys. Res. 68: 3887-3898.
- Likens, G.E. and F.H. Bormann. 1970. Chemical analyses of plant tissues from the Hubbard Brook ecosystem in New Hampshire. Yale Univ. School Forest. Bull. 79. 25p.
- List, R.J. 1966. Smithsonian meteorological tables. (6th rev. ed.) Smithsonian Misc. Collec. Vol. 114. (Publ. 4014) 527p.
- ~~Livingston~~, B.E. 1903. The distribution of the upland societies of Kent County, Michigan. Bot. Gaz. 35: 36-55.
- Linington, B.E. 1907a. Evaporation and plant development. Plant World 10: 269-276.
- Livingston, B.E. 1907b. Relative transpiration in cacti. Plant World 10: 110-114.

- Livingston, B.E. 1908a. Evaporation and centers of plant distribution. *Plant World* 11: 106-112.
- Livingston, B.E. 1908b. Evaporation and plant habitats. *Plant World* 11: 1-9.
- Livingston, B.E. and F. Shreve. 1921. The distribution of vegetation in the United States, as related to climatic conditions. Carnegie Inst. Washington, Washington, D.C. Pub. No. 284. 590p.
- Loehwing, W.F. 1951. Mineral nutrition in relation to the ontogeny of plants, pp. 343-358. *In*, Mineral nutrition of plants, E. Truog, ed. Univ. Wisconsin Press, Madison. 469p.
- Lundegårdh, H. 1951. Leaf analysis. Hilger and Watts, Ltd., London. (R.L. Mitchell, transl.) 176p.
- Macbride, T.H. 1896. Notes on forest distribution in Iowa. *Iowa Acad. Sci. Proc.* 3: 96-101.
- Mackay, K.P., Jr. 1965. An improved total hemispherical radiometer. *J. Appl. Meteorol.* 4: 112-115.
- MacMillan, C. 1892. The Metaspermae of the Minnesota valley. A list of the higher seed-producing plants indigenous to the drainage-basin of the Minnesota River. *Geol. Natur. Hist. Surv. of Minnesota, Bot. Ser. I.* 826p.
- MacMillan, C. 1899. Minnesota plant life. *Minnesota Geol. Natur. Hist. Surv., Rep. Bot. Surv.* 3. 568p.
- Mader, D. and J.A. Howarth. 1970. Relationships between foliar nutrient levels and tree growth in red pine stands in Massachusetts, pp. 205-220. *In*, Tree growth and forest soils, C.T. Youngberg and C.B. Davey, eds. (Proc. 3rd North Amer. Forest Soils Conf.). Oregon State Univ. Press, Corvallis. 527p.
- Madgwick, H.A.I. 1964. The chemical composition of foliage as an index of the nutritional status in red pine (Pinus resinosa AIT). *Plant and Soil* 21: 70-80.
- Madgwick, H.A.I. 1965. The weights and nutrient composition of understory species in ashwood. *J. Ecol.* 53: 335-341.

- Malin, J.C. 1956. The grassland of North America: its occupance and the challenge of continuous reappraisals, pp. 350-366. In, Man's role in changing the face of the earth, W.L. Thomas, Jr., ed. Univ. Chicago Press, Chicago. 1193p.
- Marks, J.B. 1942. Land use and plant succession in Coon Valley, Wisconsin. Ecol. Monogr. 12: 113-133.
- Marschner, F.J. 1930. The original forests of Minnesota. U.S.D.A. Forest Service. Map on file at North Central Forest Exp. Sta., St. Paul, Minnesota.
- Matsch, C.L. and H.E. Wright, Jr. 1967. The southern outlet of Lake Agassiz, pp. 121-140. In, Life, land and water, W.J. Mayer-Oakes, ed. (Proc. 1966 Conf. Environ. Stud. Glacial Lake Agassiz Region. Univ. Manitoba Dept. Anthropol. Occ. Paper No. 1). Univ. Manitoba Press, Winnipeg. 416p.
- McAndrews, J.H. 1964. Postglacial vegetation history of the prairie-forest transition in northwestern Minnesota. Ph.D. Thesis, Univ. Minnesota, Minneapolis. 112p.
- McAndrews, J.H. 1966. Postglacial history of prairie, savanna, and forest in northwestern Minnesota. Mem. Torrey Bot. Club 22(2): 1-72.
- McComb, A.L. and W.E. Loomis. 1944. Subclimax prairie. Bull. Torrey Bot. Club 71: 46-76.
- McCracken, H. 1959. George Catlin and the old frontier. Dial Press, New York. 216p.
- McGuire, W.W. 1834. On the prairies of Alabama. Amer. J. Sci. Arts 26: 93-98.
- McInteer, B.B. 1946. A change from grassland to forest vegetation in the "Big Barrens" of Kentucky. Amer. Midland Natur. 35: 276-282.
- McIntosh, R.P. 1967. The continuum concept of vegetation. Bot. Rev. 33: 130-187.
- Mitchell, H.L. 1936. Trends in the nitrogen, phosphorous, potassium and calcium content of the leaves of some forest trees during the growing season. Black Rock Forest Paper I(6): 30-44.

- Mitchell, H.L. and R.F. Chandler, Jr. 1939. The nitrogen nutrition and growth of certain deciduous trees of northeastern United States, with a discussion of the practice of leaf analysis as applied to forest trees. Black Rock Forest Bull. No. 11. 94p.
- Monteith, J.L. 1959a. Solarimeter for field use. J. Sci. Instruments 36: 341-346.
- Monteith, J.L. 1959b. The reflection of short-wave radiation by vegetation. Quart. J. Roy. Meteorol. Soc. 85: 386-392.
- Moseley, E.L. 1928. Flora of the oak openings west of Toledo. Ohio State Acad. Sci. Proc. (Special Paper No. 20) 8(3): 81-134.
- Moss, E.H. 1932. The vegetation of Alberta. IV. The poplar association and related vegetation of central Alberta. J. Ecol. 20: 380-415.
- Moyer, L.R. 1910a. The prairie flora of southwestern Minnesota. Minnesota Acad. Sci. Bull. 4: 357-372.
- Moyer, L.R. 1910b. The prairie legumes of western Minnesota. Minnesota Acad. Sci. Bull. 4: 373-378.
- Muir, J. 1913. The story of my boyhood and youth. Houghton Mifflin Co., New York. 294p.
- Munn, R.E. 1966. Descriptive micrometeorology. Academic Press, New York. (Adv. Geophys., Suppl. 1) 245p.
- Munter, R.C. (undated). Preparation of plant tissue for elemental analysis of Jarrell-Ash 1.5 meter spark-emission spectrometer. Unpub. Mimeo. Dept. Soil Sci., Univ. Minnesota, St. Paul. 2p.
- Murray, J.A.H., H. Bradley, W.A. Craigie, and C.T. Onions, eds. 1933 (1961). The Oxford English dictionary. Being a corrected re-issue with an introduction, supplement, and bibliography of a new English dictionary on historical principles founded mainly on the materials collected by the Philological Society. The Clarendon Press, Oxford. 12 vols., plus supplement.
- Nairn, A.E.M. 1972. Paleoclimatology: Present status. Naturwissenschaften 59: 388-393.

- Norris, D.J. and E.S. Trickett. 1968. A simple low cost pyranometer. *Solar Energy* 12: 251-253.
- Nutt, R. 1833. On the origin, extension and continuance of prairies; extracted and abridged from unpublished Mss. on a theory of the earth. *Amer. J. Sci. Arts* 23: 40-45.
- Odum, E.P. 1971. *Fundamentals of ecology*. (3rd ed.). W.B. Saunders Co., Philadelphia. 574p.
- Öpik, E.J. 1958. Climate and the changing sun. *Sci. Amer.* 198(6): 85-92.
- Ovington, J.D. 1956. The composition of tree leaves. *Forestry* 29: 22-28.
- Owensby, C.E. and K.L. Anderson. 1967. Yield responses to time of burning in the Kansas Flint Hills. *J. Range Manag.* 20: 12-16.
- Partch, M.L. 1949. Habitat studies of soil moisture in relation to plants and plant communities. Ph.D. Thesis, Univ. Wisconsin, Madison. 133p., plus appendix, bibliography, and plates.
- Partch, M.L. 1958. Vegetation changes on the Waubun prairie. *Minnesota Acad. Sci. Proc.* 25-26: 69-70.
- Partch, M.L. 1962. Species distribution in a prairie in relation to water holding capacity. *Minnesota Acad. Sci. Proc.* 30: 38-43.
- Paullin, C.O. 1932. *Atlas of the historical geography of the United States*. Carnegie Inst. Washington, Washington, D.C. Pub. No. 401. 162p. plus plates.
- Pearson, G.A. 1936. Why the prairies are treeless. *J. Forest.* 34: 405-408.
- Penfound, W.T. 1962. The savanna concept in Oklahoma. *Ecology* 43: 774-775.
- Penman, H.L. 1948. Natural evaporation from open water, bare soil, and grass. *Proc. Roy. Soc. London, Series A*, 193: 120-145.
- Penman, H.L. 1956. *Evaporation: an introductory survey*. *Neth. J. Agric. Sci.* 4: 9-29.

- Penman, H.L. 1963. Vegetation and hydrology. Commonwealth Agricultural Bureaux, Farnham Royal, Bucks, England. Tech. Commun. No. 53. 124p.
- Penman, H.L. 1967. Evaporation from forests: a comparison of theory and observation, pp. 373-380. In, Forest hydrology, W.E. Sopper and H.W. Lull, eds. Pergamon Press, New York. 813p.
- Penman, H.L., D.E. Angus, and C.H.M. van Bavel. 1967. Micro-climatic factors affecting evaporation and transpiration, pp. 483-505. In, Irrigation of agricultural lands, R.M. Hagan, H.R. Haise, and T.W. Edminster, eds. American Soc. Agron., Madison, Wisconsin. Agronomy II. 1180p.
- Pijl, L. van der. 1969. Principles of dispersal in higher plants. Springer-Verlag, Berlin (New York). 154p.
- Philip, J.R. 1959. The theory of local advection: I. J. Meteorol. 16: 535-547.
- Philip, J.R. 1961. The theory of heat flux meters. J. Geophys. Res. 66: 571-579.
- Philip, J.R. 1964. Sources and transfer processes in the air layers occupied by vegetation. J. Appl. Meteorol. 3: 390-395.
- Phillips, W.S. 1963. Photographic documentation. Vegetational changes in the northern Great Plains. Agr. Exp. Sta., Coll. Agr., Univ. Arizona, Tucson. Rep. 214. 185p.
- Planck, M. 1914 (1959). The theory of heat radiation. Dover Publ., Inc., New York. 224p.
- Plass, G.N. 1959. Carbon dioxide and climate. Sci. Amer. 201(1): 41-47.
- Platt, R.B. and J.F. Griffiths. 1964. Environmental measurement and interpretation. Reinhold Publ. Corp., New York. 235p.
- Plummer, F.G. 1912a. Forest fires: their causes, extent and effects, with a summary of recorded destruction and loss. U.S.D.A. Forest Service. Bull. No. 117. 39p.
- Plummer, F.G. 1912b. Lightning in relation to forest fires. U.S.D.A. Forest Service. Bull. No. 111. 39p.

- Pluth, D.J., R.S. Adams, Jr., R.H. Rust, and J.R. Peterson. 1970. Characteristics of selected horizons from 16 soil series in Minnesota. Univ. Minnesota Agr. Exp. Sta. Tech. Bull. 272. 34p.
- Pluth, D.J. and H.F. Arneman. 1965. Forest soil and tree growth characteristics related to a synecological coordinate system, pp. 331-351. In, Forest-soil relationships in North America, C.T. Youngberg, ed. (Papers presented at Second North American Forest Soils Conf.) Oregon State Univ. Press, Corvallis. 532p.
- Pool, R.J. 1915. The invasion of a planted prairie grove. Soc. Amer. Forest., Proc. 10: 1-8.
- Pool, R.J., J.E. Weaver, and F.C. Jean. 1918. Further studies in the ecotone between prairie and woodland. Nebraska Univ., Univ. Stud. 18: 7-53.
- Pound, R. and F.E. Clements. 1898. The vegetation regions of the prairie province. Bot. Gaz. 25: 381-394.
- Radiation Commission of IAM. 1958. IGY instruction manual, part IV. Radiation instruments and measurements. Ann. Geophys. Year V: 363-466.
- Raupach, M., R. Boardman, and A.R.P. Clarke. 1969. Growth rates of Pinus radiata D. Don. in relation to foliar levels of nitrogen and phosphorus for plantations in the southeast of south Australia. C.S.I.R.O., Australia. Soil Pub. No. 26. 28p.
- Record, F.A. and H.E. Cramer. 1966. Turbulent energy dissipation rates and exchange processes above a non-homogeneous surface. Quart. J. Roy. Meteorol. Soc. 92: 519-532.
- Redway, J.W. 1894. The treeless plains of the United States. Geogr. J. 3: 209-213.
- Reifsnyder, W.E. and H.W. Lull. 1965. Radiant energy in relation to forests. U.S.D.A. Forest Service. Tech. Bull. No. 1344. 111p.
- Rider, N.E., J.R. Philip, and E.F. Bradley. 1963. The horizontal transport of heat and moisture--a micrometeorologic study. Quart. J. Roy. Meteorol. Soc. 89: 507-531.
- Robinson, N. 1966. Solar radiation. Elsevier Publ. Co., New York. 347p.

- Robinson, N. and L. Stoch. 1964. Sky radiation measurements and corrections. *J. Appl. Meteorol.* 3: 179-181.
- Rose, C.W. 1966. *Agricultural physics*. Pergamon Press, New York. 226p.
- Rosenberg, N.J. and K.W. Brown. 1973. Measured and modelled effects of microclimate modification on evapotranspiration by irrigated crops in a region of strong sensible heat advection, pp. 539-546. In, Plant response to climatic factors, R.O. Slatyer, ed. UNESCO, Ecology and Conservation, 5. (Proc. Uppsala Symp., 1970). 574p.
- Rosendahl, C.O. 1948. A contribution to the knowledge of the Pleistocene flora of Minnesota. *Ecology* 29: 284-315.
- Rosendahl, C.O. 1955 (1928). *Trees and shrubs of the Upper Midwest*. Univ. Minnesota Press, Minneapolis. 411p.
- Rowe, J.S. 1956. Uses of undergrowth plant species in forestry. *Ecology* 37: 461-473.
- Russell, R.J. 1932. Dry climates of the United States. II. Frequency of dry and desert years 1901-1920. Univ. California Pub. Geogr. 5: 245-274.
- Russell, R.J. 1956. Environmental changes through forces independent of man, pp. 453-470. In, Man's role in changing the face of the earth, W.L. Thomas, Jr., ed. Univ. Chicago Press, Chicago. 1193p.
- Rutter, A.J. 1968. Water consumption by forest, pp. 23-84. In, Water deficits and plant growth, Vol. II. Plant water consumption and response, T.T. Kozlowski, ed. Academic Press, New York. 333p.
- Sargent, C.S. 1884. Report on the forests of North America (exclusive of Mexico). Dept. Interior, U.S. Census Office, 10th Census of the United States, Vol. 9. 612p.
- Sauer, C.O. 1950. Grassland climax, fire, and man. *J. Range Manag.* 3: 16-21.
- Sauer, C.O. 1956. The agency of man on the earth, pp. 49-69. In, Man's role in changing the face of the earth, W.L. Thomas, Jr., ed. Univ. Chicago Press, Chicago. 1193p.

- Schell, I.I. 1961. Recent evidence about the nature of climate changes and its implications. New York Acad. Sci., Ann. 95: 251-270.
- Schimper, A.F.W. 1903. Plant-geography upon a physiological basis. Clarendon Press, Oxford. 839p.
- Scholte Ubing, D.W. 1959. Over stralingsmetingen, de warmtebalans en de verdamping van gras. (Studies on solar and net radiation and on evapotranspiration of grass). Mededelingen van de Landbouwhogeschool te Wageningen, Nederland 59(10): 1-93.
- Schwartz, G.M. and G.A. Thiel. 1954. Minnesota's rocks and waters. A geologic survey. Minnesota Geol. Surv. Bull. 37. 366p.
- Schopmeyer, C.S. 1961. Foliar absorption and translocation of phosphorus by loblolly pine seedlings. Forest Sci. 7: 330-336.
- Scott, D.R.M. 1955. Amount and chemical composition of the organic matter contributed by overstory and understory vegetation to forest soil. Yale Univ. School Forest. Bull. 62. 73p., plus illustrations.
- Seal, H.L. 1964. Multivariate statistical analysis for biologist. Methuen and Co. Ltd., London. 209p.
- Sears, P.B. 1932. Postglacial climate in eastern North America. Ecology 13: 1-6.
- Sears, P.B. 1948. Forest sequence and climatic change in northeastern North America since early Wisconsin time. Ecology 29: 326-333.
- Sellers, W.D. 1965. Physical climatology. Univ. Chicago Press, Chicago. 272p.
- Shaler, N.W. 1891. The origin and nature of soils, pp. 213-345. In, Twelfth Annual Report of the U.S. Geological Survey, J.W. Powell, Dir. Part 1 - Geology. 675p.
- Shantz, H.L. 1954. The place of grasslands in the earth's cover of vegetation. Ecology 35: 143-145.
- Shantz, H.L. and R. Zon. 1924. Natural vegetation, 29p. In, Atlas of American Agriculture (1936). U.S. Dept. Agr., Washington, D.C.

Shapley, H., ed. 1953. Climatic change. Evidence, causes, and effects. Harvard Univ. Press, Cambridge, Massachusetts. 318p.

Shay, C.T. 1965. Postglacial vegetational development in northwestern Minnesota and its implications for pre-historic human ecology. M.A. Thesis, Univ. Minnesota, Minneapolis. 112p.

Shay, C.T. 1967. Vegetation history of the southern Lake Agassiz basin during the past 12,000 years, pp. 231-252. In, Life, land and water, W.J. Mayer-Oakes, ed. (Proc. 1966 Conf. Environ. Stud. Glacial Lake Agassiz Region. Univ. Manitoba Dept. Anthropol. Occ. Paper No. 1). Univ. Manitoba Press, Winnipeg. 416p.

Shimek, B. 1900. The distribution of forest trees in Iowa. Iowa Acad. Sci. Proc. 7: 47-59.

Shimek, B.A. 1910. Prairie openings in the forest. Iowa Acad. Sci. Proc. 17: 16-19.

Shimek, B. 1911. The prairies. Bull. Labs. Natur. Hist. Univ. Iowa 6(3): 169-240, plus plates.

Shimek, B.A. 1924. Prairies of the Mississippi bluffs. Iowa Acad. Sci. Proc. 31: 205-212.

Shirley, H.L. 1936. Will pine or aspen dominate Minnesota forests? Minnesota Acad. Sci. Proc. 4: 25-28.

Shul'gin, A.M. 1957 (1965). The temperature regime of soils. GIMIZ. Gidrometeorologicheskoe Izdatel'stvo. Leningrad. Transl. TT65-50083. Office of Tech. Serv., Washington, D.C. 218p.

Siccama, T.G., F.H. Bormann, and G.E. Likens. 1970. The Hubbard Brook ecosystem study: Productivity, nutrients, and phytosociology of the herbaceous layer. Ecol. Monogr. 40: 389-402.

Simonsen, R.W. 1941. Studies of buried soils formed from till in Iowa. Soil Sci. Soc. Amer. Proc. 6: 373-381.

Singer, S.F., ed. 1970. Global effects of environmental pollution. Springer-Verlag, New York. 218p.

Slatyer, R.O. 1967. Plant-water relationships. Academic Press, New York. 366p.

Slatyer, R.O. and I.C. McIlroy. 1961. Practical microclimatology; with special reference to the water factor in soil-plant-atmosphere relationships. CSIRO, Australia. (UNESCO). 211p., plus figures and appendices.

Smeins, F.E. and D.E. Olsen. 1970. Species composition and production of a native northwestern Minnesota tall grass prairies. Amer. Midland Natur. 84: 398-410.

Smith, E.F. and C.E. Owensby. 1973. Effects of fire on true prairie grasslands. Proc. Ann. Tall Timbers Fire Ecology Conf. 12: 9-22.

Smith, G.D., W.H. Allaway, and F.F. Riecken. 1950. Prairie soils of the Upper Mississippi Valley. Adv. Agron. 2: 157-205.

Smith, P.F. 1962. Mineral analysis of plant tissues. Ann. Rev. Plant Physiol. 13: 81-108.

Smith, W.O. 1940. Thermal conductivities in moist soils. Soil Sci. Soc. Amer. Proc. (1939) 4: 32-40.

Smith, W.O. and H.G. Byers. 1939. The thermal conductivity of dry soils of certain great soil groups. Soil Sci. Soc. Amer. Proc. (1938) 3: 13-19.

Snedecor, G.W. 1946 (1950). Statistical methods applied to experiments in agriculture and biology. Iowa State College Press, Ames. 485p.

Soil Science Department, University of Minnesota, Soil Conservation Service, U.S. Department of Agriculture, and Minnesota Geological Survey. 1969. Minnesota soil atlas. Brainerd sheet. Univ. Minnesota Agr. Exp. Sta. Misc. Rep. 90. 35p., plus map.

Soil Science Department, University of Minnesota, Soil Conservation Service, U.S. Department of Agriculture, and Minnesota Geological Survey. 1971. Minnesota soil atlas. Hibbing sheet. Univ. Minnesota Agr. Exp. Sta. Misc. Rep. 110. 47p., plus map.

Spurr, S.H. 1964. Forest ecology. Ronald Press, New York, 352p.

Stallard, H. 1929. Secondary succession in the climax forest formations of northern Minnesota. Ecology 10: 476-547.

- Steel, R.G.D. and J.H. Torrie. 1960. Principles and procedures of statistics with special reference to the biological sciences. McGraw-Hill Book Co., Inc. New York. 481p.
- Steiger, T.L. 1930. Structure of prairie vegetation. Ecology 11: 170-217.
- Stewart, O.C. 1951. Burning and natural vegetation in the United States. Geogr. Rev. 41: 317-320.
- Stewart, O.C. 1953. Why the Great Plains are treeless. The Colorado Quart. 2(1): 40-50.
- Stewart, O.C. 1956. Fire as the first great force employed by man, pp. 115-133. In, Man's role in changing the face of the earth, W.L. Thomas, Jr., ed. Univ. Chicago Press, Chicago. 1193p.
- Stout, A.B. 1944. The bur oak openings in southern Wisconsin. Wisconsin Acad. Sci. Arts Letters, Trans. 36: 141-161.
- Sukachev, V. and N. Dylis. 1964. Fundamentals of forest biogeocoenology. Oliver and Boyd, London. 672p.
- Suomi, V.E. and P.M. Kuhn. 1958. An economical net radiometer. Tellus 10: 160-163.
- Suomi, V.E. and C.B. Tanner. 1958. Evapotranspiration estimates from heat-budget measurements over a field crop. Amer. Geophys. Union Trans. 39: 298-304.
- Swan, H.S.D. 1962. The mineral nutrition of the Grand'Mere plantations. Pulp and Paper Res. Inst. Can. Tech. Rep. No. 276. (Woodlands Res. Index 131). 14p.
- Swinbank, W.C. 1958. Turbulent transfer in the lower atmosphere, pp. 35-37. In, Climatology and microclimatology. (Proc. Canberra Symposium). UNESCO. Arid Zone Research 11. 355p.
- Szeicz, G. 1965. A miniature tube solarimeter. J. Appl. Ecol. 2: 145-147.
- Szeicz, G. 1966. Field measurements of energy in the 0.4-0.7 micron range, pp. 41-51. In, Light as an ecological factor, R. Bainbridge, G.C. Evans, and O. Rackham, eds. (British Ecological Symposium Vol. 6). (Blackwell Scientific Publ.). John Wiley and Sons, Inc., New York. 452p.

- Szeicz, G., J.L. Monteith, and J.M. dos Santos. 1964. Tube solarimeter to measure radiation among plants. *J. Appl. Ecol.* 1: 169-174.
- Tajchman, S.J. 1972. The radiation and energy balances of coniferous and deciduous forests. *J. Appl. Ecol.* 9: 359-375.
- Tamm, C.O. 1964. Determination of nutrient requirements of forest stands. *Int. Rev. Forest Res.* 1: 115-170.
- Tanner, C.B. 1957. Factors affecting evaporation from plants and soils. *J. Soil and Water Conserv.* 12: 221-227.
- Tanner, C.B. 1960a. A simple aero-heat budget method for determining daily evapotranspiration. *Int. Congr. Soil Sci. Trans.* 7, Madison, I: 203-209.
- Tanner, C.B. 1960b. Energy balance approach to evapotranspiration from crops. *Soil Sci. Soc. Amer. Proc.* 24: 1-9.
- Tanner, C.B. 1967. Measurement of evapotranspiration, pp. 534-574. *In*, Irrigation of agricultural lands, R.M. Hagen, H.R. Haise, and T.W. Edminster, eds. *Amer. Soc. Agron.*, Madison, Wisconsin. Agronomy II. 1180p.
- Tanner, C.B. 1968. Evaporation of water from plants and soil, pp. 73-106. *In*, Water deficits and plant growth, Vol. 1. Development, control, and measurement, T.T. Kozlowski, ed. Academic Press, New York. 390p.
- Tanner, C.B., C.A. Federer, T.A. Black, and J.B. Swan. 1969. Economical radiometer theory, performance and construction. College of Agricultural and Life Sciences, Univ. Wisconsin, Madison. Res. Rep. 40. 86p.
- Tanner, C.B. and M. Fuchs. 1968. Evaporation from unsaturated surfaces: a generalized combination method. *J. Geophys. Res.* 73: 1299-1304.
- Tanner, C.N. and E.R. Lemon. 1962. Radiant energy utilized in evapotranspiration. *Agron. J.* 54: 207-212.
- Tester, J.R. 1965. Effects of a controlled burn on small mammals in a Minnesota oak-savanna. *Amer. Midland Natur.* 74: 240-243.

- Tester, J.R. and W.H. Marshall. 1962. Minnesota prairie management techniques and their wildlife implications. North Amer. Wildlife and Natur. Res. Conf. Trans. 27: 267-287.
- Thompson, J.R. and A.D. Ozment. 1972. The Rocky Mountain millivolt integrator for use with solar radiation sensors. U.S.D.A. Forest Service, Rocky Mountain Forest and Range Exp. Sta. Res. Note RM - 225. 8p.
- Thomson, J.W., Jr. 1940. Relic prairie areas in central Wisconsin. Ecol. Monogr. 10: 685-717.
- Thurtell, G.W. and C.B. Tanner. 1964. Electronic integrator for micrometeorological data. J. Appl. Meteorol. 3: 198-202.
- Todd, J.E. 1878. Notes on the distribution of timber in south-western Iowa, with inferences concerning the origin of prairies. Amer. Natur. 12: 91-96.
- Transeau, E.N. 1905. Forest centers of eastern America. Amer. Natur. 39: 875-889.
- Transeau, E.N. 1935. The prairie peninsula. Ecology 16: 423-437.
- Truman, H.V. 1937. Fossil evidence of two prairie invasions of Wisconsin. Wisconsin Acad. Sci. Arts Letters, Trans. 30: 35-42.
- Upham, W. 1880. V. Preliminary report on the geology of central and western Minnesota, pp. 70-125. In, The geological and natural history survey of Minnesota, 8th Ann. Rep. (1879), N.H. Winchell, ed. Pioneer Press Co., St. Paul. 187p.
- Upham, W. 1884. Catalogue of the flora of Minnesota, including its phaenogamous and vascular cryptogamous plants, indigenous, naturalized, and adventive. In, The geological and natural history survey of Minnesota, 12th Ann. Rep. (1883), Part VI, N.H. Winchell, ed. Johnson, Smith, and Harrison, Minneapolis. 193p.
- Upham, W. 1892. Geographic limits of species of plants in the basin of the Red River of the North. Boston Soc. Natur. Hist. Proc. 25: 140-172.

Upham, W. 1895 (1896). The Glacial Lake Agassiz. Monogr.
U.S. Geol. Surv., Vol. 25. 658p.

U.S. Dept. Commerce. National Oceanic and Atmospheric Administration (Weather Bureau; Environmental Sciences Services Administration). 1961-71a. Climatological data. Annual summary. Minnesota [Vols. 67-77(13)]; North Dakota [Vols. 70-80(13)]; South Dakota [Vols. 66-76(13)]. Washington, D.C.

U.S. Dept. Commerce. National Oceanic and Atmospheric Administration (Weather Bureau; Environmental Sciences Services Administration). 1961-71b. Climatological data. National summary. Washington, D.C. Vol. 12-22(1-13).

U.S. Dept. Commerce. National Oceanic and Atmospheric Administration. 1971. Daily weather maps, weekly series (July 5-11; July 19-25; August 2-8). U.S. Dept. Commerce, NOAA. Washington, D.C. 8p.(ea.).

U.S. Dept. Commerce. National Oceanic and Atmospheric Administration. 1972a. PERSUM. Normals program for cooperating stations. Preliminary listing of average monthly temperature for 1941-1970. State #21. Minnesota. Environmental Data Service, National Climatic Center, Asheville, North Carolina. 74p.

U.S. Dept. Commerce. National Oceanic and Atmospheric Administration. 1972b. PERSUM. Normals program for cooperating stations. Preliminary listing of total monthly precipitation for 1941-1970. State #21. Minnesota. Environmental Data Service, National Climatic Center, Asheville, North Carolina. 86p.

Van Bavel, C.H.M. 1966. Potential evaporation: the combination concept and its experimental verification. Water Resources Res. 2: 455-467.

Van Wijk, W.R. 1963. General temperature variations in a homogeneous soil, pp. 144-170. In, Physics of the plant environment, W.R. Van Wijk, ed. John Wiley and Sons, Inc., New York. 382p.

Van Wijk, W.R. and W.J. Derksen. 1966. Thermal properties of a soil near the surface. Agr. Meteorol. 3: 333-342.

Van Wijk, W.R. and D.W. Scholte Ubing. 1963. Radiation, pp. 62-101. In, Physics of the plant environment, W.R. Van Wijk, ed. North-Holland Publ. Co., Amsterdam. 382p.

- Vestal, A.G. 1918a. Invasion of forest land by prairie along railroads. Illinois State Acad. Sci., Trans. 11: 126-128.
- Vestal, A.G. 1918b. Local inclusions of prairie within forest. Illinois State Acad. Sci., Trans. 11: 122-126.
- Viro, P.J. 1961. Evaluation of site fertility. Unasylva 15: 91-97.
- Voss, J. 1934. Postglacial migration of forests in Illinois, Wisconsin, and Minnesota. Bot. Gaz. 96: 3-43.
- Wadleigh, C.H. and L.A. Richards. 1951. Soil moisture and the mineral nutrition of plants, pp. 411-450. In, Mineral nutrition of plants, E. Truog, ed. Univ. Wisconsin Press, Madison. 469p.
- Waring, R.H. and J. Major. 1964. Some vegetation of the California redwood region in relation to gradients of moisture, nutrients, light, and temperature. Ecol. Monogr. 34: 167-215.
- Warming, E. 1909. Oecology of plants. An introduction to the study of plant-communities. Clarendon Press, Oxford. 422p.
- Warnes, D.D., S.D. Evans, and E.A. Hallauer. 1971. Eighty-four years of weather reporting in west central Minnesota. Part 1. Averages and extremes. Univ. Minnesota Agr. Exp. Sta. Misc. Rep. 108. 55p.
- Watt, R.F. and M.L. Heinzelman. 1965. Foliar nitrogen and phosphorus level related to site quality in a northern Minnesota spruce bog. Ecology 46: 357-361.
- Watts, W.A. and R.C. Bright. 1968. Pollen, seed, and mollusk analysis of a sediment core from Pickeral Lake, Day County, South Dakota. Bull. Geol. Soc. Amer. 79: 855-876.
- Weaver, J.E. and F.W. Albertson. 1956. Grasslands of the Great Plains. Their nature and use. Johnsen Publ. Co., Lincoln, Nebraska. 395p.
- Weaver, J.E. and F.E. Clements. 1938. Plant ecology. 2nd ed. McGraw-Hill Book Co., New York. 601p.
- Weaver, J.E. and T.J. Fitzpatrick. 1934. The prairie. Ecol. Monogr. 4: 109-295.

- Weaver, J.E. and J. Kramer. 1932. Root system of Quercus macrocarpa in relation to the invasion of prairie. Bot. Gaz. 94: 51-85.
- Weaver, J.E. and A.F. Thiel. 1917. Ecological studies in the tension zone between prairie and woodland. Bot. Surv. Nebraska (Univ. Nebraska), New Ser., No. 1. 60p.
- Wells, P.V. 1965. Scarp woodlands, transported grassland soils, and concept of grassland climate in the Great Plains region. Science 148: 246-249.
- Wells, P.V. 1970. Historical factors controlling vegetation patterns and floristic distributions in the Central Plains region of North America, pp. 211-221. In, Pleistocene and Recent environments of the Central Great Plains, W. Dort, Jr. and J.K. Jones, Jr., eds. Dept. Geology, Univ. Kansas Spec. Pub. 3. Univ. Kansas Press, Lawrence. 433p.
- Wells, R.W. 1819. On the origin of prairies. Amer. J. Sci. more especially of mineralogy, geology, and the other branches of natural history; including agriculture and the ornamental as well as useful arts (1st Ser., 2nd ed.) 1: 331-337.
- White, C.A. 1868. The lakes of Iowa, --past and present. Amer. Natur. 2: 143-155.
- White, C.A. 1871. Prairie fires. Amer. Natur. 5: 68-70.
- White, D.P. 1954. Variation in the nitrogen, phosphorus, and potassium contents of pine needles with season, crown position, and sample treatment. Soil Sci. Soc. Amer. Proc. 18: 326-330.
- White, E.H. and M.C. Carter. 1970. Relationships between foliage nutrient levels and growth of young natural stands of Populus deltoides Bartr., pp. 283-294. In, Tree growth and forest soils, C.T. Youngberg and C.B. Davey, eds. (Proc. 3rd North Amer. Forest Soils Conf.). Oregon State Univ. Press, Corvallis. 527p.
- Whitford, P.B. 1958. Species distribution and age of hardwood stands in the prairie forest region of Wisconsin. Ph.D. Thesis, Univ. Wisconsin, Madison. 111p., plus bibliography and appendices.
- Whitford, P.B. 1958. A study of prairie remnants in southeastern Wisconsin. Ecology 39: 727-733.

Whitford, P.B. and K. Whitford. 1971. Savanna in central Wisconsin, U.S.A. *Vegetatio* 23: 77-87.

Whitman, A. 1877. VII. Entomology, pp. 128-161. In, The geological and natural history survey of Minnesota, 5th Ann. Rep. (1876), N.H. Winchell, ed. 248p.

Whitman, A. 1878. X. Entomology, pp. 132-153. In, The geological and natural history survey of Minnesota, 6th Ann. Rep. (1877), N.H. Winchell, ed. 226p.

Whitney, J.D. 1876a. Plain, prairie, and forest. Part 1. *Amer. Natur.* 10: 577-588.

Whitney, J.D. 1876b. Plain, prairie, and forest. Part 2. *Amer. Natur.* 10: 656-667.

Wilde, S.A. 1954. Mycorrhizal fungi: their distribution and effect on tree growth. *Soil Sci.* 78: 23-31.

Wilde, S.A. 1958. Diagnosis of nutrient deficiencies by foliar and soil analysis in silvicultural practice, pp. 138-140. In, First North American forest soils conference, T.D. Stevens and R.L. Cook, eds. *Agr. Exp. Sta., Michigan State Univ., East Lansing.* 226p.

Wilde, S.A., G.K. Voigt, and J.G. Iyer. 1964. Soil and plant analysis for tree culture. Oxford Publ. House, Calcutta, India. 209p.

Will, G.F. 1946. Tree ring studies in North Dakota. *North Dakota Agr. Exp. Sta. Bull.* 338. 24p.

Willard, D.E. 1922. The story of the North Star State. Webb Publ. Co., St. Paul, Minnesota. 395p.

Willetts, H.C. 1949. Long-period fluctuations of the general circulation of the atmosphere. *J. Meteorol.* 6: 34-50.

Willetts, H.C. 1961. The pattern of solar climatic relationships. *New York Acad. Sci., Ann.* 95: 89-106.

Wilson, W.H. and T.D. Epps. 1920. The construction of thermocouples by electro-deposition. *Phys. Soc. London, Proc.* 32: 326-340.

Winchell, A. 1864. On the origin of the prairies of the valley of the Mississippi. *Amer. J. Sci. Arts, Second Ser.* 38: 332-344.

- Winchell, N.H. 1875. Notes on the Big Woods. Minnesota State Hort. Soc. Trans. (1875): 47-50.
- Winchell, N.H. and S.F. Peckham. 1874. The geological and natural history survey of Minnesota. Second Ann. Rep. (1873). 148p.
- Withrow, R.B. 1951. Light as a modifying influence on the mineral nutrition of plants, pp. 389-410. In, Mineral nutrition of plants, E. Truog, ed. Univ. Wisconsin Press, Madison. 469p.
- Woodard, J. 1924. Origin of prairies in Illinois. Bot. Gaz. 77: 241-261.
- World Data Center A for Solar-Terrestrial Physics. 1973. Zurich final relative sunspot numbers for 1971 and 1972. National Oceanic and Atmospheric Administration, Boulder Colorado.
- Wright, H.E., Jr. 1964a. Aspects of the early postglacial forest succession in the Great Lakes region. Ecology 45: 439-448.
- Wright, H.E., Jr. 1964b. The classification of the Wisconsin glacial stage. J. Geol. 72: 628-637.
- Wright, H.E., Jr. 1968. History of the prairie peninsula, pp. 78-88. In, The Quaternary of Illinois, R.E. Bergstrom, ed. Univ. Illinois Coll. Agr. Spec. Pub. 14. 179p.
- Wright, H.E., Jr. 1970. Vegetational history of the central plains, pp. 157-172. In, Pleistocene and Recent environments of the Central Great Plains, W. Dort, Jr., and J.K. Jones, Jr., eds. Dept. Geology, Univ. Kansas Spec. Pub. 3. Univ. Kansas Press, Lawrence. 433p.
- Wright, H.E., Jr. 1971. Late Quaternary vegetational history of North America, pp. 425-464. In, The Late Cenozoic Glacial Ages, K.K. Turekian, ed. Yale Univ. Press. 606p.
- Wright, H.E., Jr. and R.V. Ruhe. 1965. Glaciation of Minnesota and Iowa, pp. 29-41. In, The Quaternary of the United States, H.E. Wright, Jr. and D.G. Frey, eds. Princeton Univ. Press, Princeton. 922p.
- Wright, H.E., Jr. and W.A. Watts with contributions by S. Jelgersma, J.C.B. Waddington, J. Ogawa, and T.C. Winter. 1969. Glacial and vegetational history of northeastern Minnesota. Minnesota Geol. Surv. Spec. Pub. 11. 59p.

- Wright, H.E. Jr., T.C. Winter, and H.L. Patten. 1963. Two pollen diagrams from southeastern Minnesota: Problems in the regional late-glacial and postglacial vegetational history. Geol. Soc. Amer. Bull. 74: 1371-1396.
- Youngberg, C.T. 1958. The uptake of nutrients by western conifers in forest nurseries. J. Forest. 56: 337-340.
- Zedler, J. and O.L. Loucks. 1969. Differential burning response of Poa pratensis fields and Andropogon scoparius prairies in central Wisconsin. Amer. Midland Natur. 81: 341-352.
- Zon, R. and W.N. Sparhawk. 1923. Forest resources of the world. McGraw-Hill Book Co., Inc., New York. 2 Vols. 997p.

5.0 APPENDIX

5.1 Study Area Descriptions

(Legal descriptions are based on the 5th Principal Meridian.)

Stand 1. Location: Winona Co., 5.5 mi. N of St. Charles, Whitewater State Park, Chimney Rock Bluff; NE $\frac{1}{4}$ SE $\frac{1}{4}$ Sec. 20, T 107 N, R 10 W. Topography: Steep hills, unglaciated, N 30° W aspect, 0-20% slope, ridgetop, rock outcrops, rocky soil. Vegetation: 60-70% canopy dominated by Juniperus virginiana (2-10" D.B.H.), Quercus alba, Q. borealis (0-14"), Ostrya virginiana, with grasses dominant in the understory; bare soil 10% excluding trails. Remarks: Maple-basswood climax vegetation on moist side slope. Survey records: April, 1834; at SE corner of Sec. 20, white oak (2", 10"), basswood (8"), and aspen (7") recorded as witnesses; at half section post between sections 20 and 21, white oak (8") and black oak (6") recorded; at NE corner of Sec. 20, bur oak (10", 12"), white birch (5"), and aspen (6") recorded; comments around the south and east sides of Sec. 20 include: "Land broken," "2nd [and] 3rd rate" (marginal or not suitable for agriculture), "very little timber," some in groves, hazel and shrub oak undergrowth, dead aspen on hillside.

STAND 2. Location: Jackson Co., 7 mi. N and 4 mi. W of Jackson, Kilen Woods State Park; W $\frac{1}{2}$ NE $\frac{1}{4}$ Sec. 17, N $\frac{1}{4}$ SW $\frac{1}{4}$ Sec. 16, T 103 N, R 35 W. Topography: Mostly level land intersected by large draw with perennial stream; N aspect, 0-5% slope, upland above draw, good soil. Vegetation: 60% Quercus macrocarpa (1-20") with Ulmus americana (6-28") and Tilia americana; bur oak and basswood reproduction; much

Ribes americanum; no bare soil. Remarks: stand not open grown. Survey records: October, 1858; 23.00 ch. east between Sections 16 and 21 scattered bur oak bears NW and SE; at half section between 16 and 21, white oak (24"), bur oak (8") recorded; north between Sections 16 and 17, 15.00 ch. enter timber, 35.00 ch. leave timber, 40.00 ch. prairie vegetation; comments at the southeast corner of Sec. 17 include "surface rolling, prairie & timber, soil 1st rate. Timber mostly bur oak." North between Sections 16 and 17, "Surface rolling, prairie & timber, soil 1st rate;" oak, elm, ash, basswood, ironwood; between Sections 17 and 8, "Surface rolling, prairie, soil 2nd rate."

STAND 3. Location: Rock Co., 5 mi. N of Luverne, Blue Mounds State Park; NE $\frac{1}{4}$ NW $\frac{1}{4}$ Sec. 24, T 103 N, R 45 W. Topography: Rolling surface with rock outcrops, S 80° E aspect, 0-30% slope, creek bottom and adjacent side slope, rocky and sandy soil. Vegetation: 40% canopy of Fraxinus pennsylvanica (1-15") and Quercus macrocarpa (1-12"), oak reproduction, dense understory, primarily riparian habitat; 15% exposed rock. Remarks: Grazed in the past, now occasionally mowed; water impoundments on stream; some trees planted. Survey records: November, 1869; between Sections 13 and 24, 14 and 23, and 23 and 24, prairie land; comments include "Stony prairie," "slough," "prairie land;" no mention of trees. [note: subsequent reexamination of the area in another part of the park

showed mesic forest invading below a bluff and Rhus glabra and bur oak invading prairie on the top of the bluff.]

STAND 4. Location: Murray Co., 11 mi. NE of Slayton, Shetek State Park; SE $\frac{1}{4}$ NE $\frac{1}{4}$ Sec. 6, T 107 N, R 40 W. Topography: Rolling to level surface on bench above Lake Shetek, west aspect, 0-5% slope, sandy-gravelly soil. Vegetation: 40-50% canopy of mature Quercus macrocarpa (up to 36") and Fraxinus pennsylvanica (up to 28") with Ulmus americana and Acer negundo; shrub understory; no bare soil. Remarks: Oaks open grown; good reproduction of species other than oaks; herbaceous ground cover dense, perhaps the result of disturbance. Survey records: September, 1861; between Sections 6 and 7, "grove of scattering Br. Oak" and prairie vegetation; north between Sections 5 and 6 11.50 ch. "leave grove of shrubby bur oak; enter prairie," 19.00 ch., "small grove on slough," 56.50 ch. "hay marsh," 70.00 ch. "scattering of young timber," 74.00 ch. "brush," 76.00 ch. "cane slough;" 11.60 ch. north of southeast corner of sec. 6., bur oak (4", 10").

STAND 5. Location: Lyon Co., 14 mi. S of Marshall; NW $\frac{1}{4}$ SW $\frac{1}{4}$ Sec. 11, T 109 N, R 40 W. Topography: Rolling with draws and small seasonal streams; uplands above Cottonwood River; S 60° E aspect, 5-10% slope, upper slope. Vegetation: 50% canopy of Quercus macrocarpa (10-15"), shrubby understory of Xanthoxylum americanum, Symphoricarpos occidentalis, and

grasses; no bare soil. Remarks: Has been grazed in the past but not for at least 5 years; bur oaks do not have open grown appearance. Survey records: July-August, 1867; SW corner of Sec. 11, prairie; no trees mentioned between Sections 10 and 11; comments include "Surface rolling" and "Soil No. 1."

STAND 6. Location: Pope Co., 3 mi. NW of Glenwood; NE $\frac{1}{4}$ SE $\frac{1}{4}$ Sec. 28, T 126 N, R 38 W. Topography: Top of rolling hills, above marshland and lakes; N 70° E aspect, 0-5% slope, sand to loam soil. Vegetation: 60-70% canopy of Quercus macrocarpa (up to 12"), with understory of Acer negundo and Xanthoxylum americanum with good herbaceous ground cover; no bare soil (at least litter covered). Remarks: Stand cut over 20-30 years ago, since then not much disturbance; oak stump sprouts, some oak reproduction. Survey records: September, 1859; SE corner of Sec. 28, bur oak (6", 6", 6", 24"); north between Sections 27 and 28, 38.00 ch. "enter bur oak opening," 40.00 ch., bur oak (20", 20"), 42.50-56.55 ch. "wet marsh;" S edge of Sec. 28, 40.00 ch., prairie; comments include "Surface rolling" with prairie and "timber bur oak," "soil 1st rate" and "Surface hilly, prairie soil 2nd rate."

STAND 7. Anoka Co., East Bethel, Allison Savanna Nature Conservancy Area; NE $\frac{1}{4}$ NE $\frac{1}{4}$ Sec. 2, T 133 N, R 23 W. Topography: Rolling sand dunes; south aspect, 0-5% slope, upper slope. Vegetation: Oak savanna with Quercus macrocarpa and

Q. borealis dominant; understory predominantly prairie with occasional Corylus americana; 5% bare soil. Remarks: Some controlled burning in area, not in study plot area; gopher mounds and ant hills common; some blowouts and marshy areas. Survey records: November, 1856; NE corner of Sec. 2, black oak (11", 11"); south between Sections 1 and 2 at 41.50 ch. black oak (10"); comments include "Sand rolling," "3rd rate soil;" "Scattering black oak & bur oak;" "tamarack, and oak brush."

STAND 8, 9. Ottertail Co., $3\frac{1}{2}$ mi. W of Urbank, Inspiration Peak State Park; $SE\frac{1}{4}$ $SW\frac{1}{4}$ Sec. 22 and $NE\frac{1}{4}$ $NW\frac{1}{4}$ Sec. 27, T 131 N, R 39 W. Topography: Hill above generally rolling country; hilltop and N 60° W aspect sidehill stands, slope 0%, 30%; gravelly soil. Vegetation: Stand 8 (hilltop)—Open stand of Quercus macrocarpa with shrub and prairie understory; bare soil 5%. Stand 9 (sideslope)—Closed stand of Quercus macrocarpa, Q. ellipsoidalis, and Ulmus americana with shrubby and herbaceous understory; no bare soil. Remarks: Hilltop dry, sideslope moist; the area was probably grazed in the past. Survey records: July, 1871; W between Sections 22 and 27, 40.10 ch. oak (14", 16"); SW corner of Sec. 22, oak (4", 6", 6", 6"); comments include "Surface very rolling;" "Scattered oak timber;" "Thick oak brush;" "Open brush land;" "Soil second rate."

STAND 10. Location: Hubbard Co., $3/4$ mi. S of Hubbard;

W $\frac{1}{2}$ SE $\frac{1}{4}$ SW $\frac{1}{4}$ Sec. 29, T 139 N, R 34 W. Topography: Level land above slightly rolling topography; sandy soil. Vegetation: 70% canopy cover of Quercus macrocarpa (4-6", 25-30' tall), with a few Pinus banksiana; well developed shrub layer; no bare soil. Remarks: No visible results of fire although some oaks clumped at root collar; a few large open grown oaks present, but most of canopy was closed grown. Survey records: December, 1860; SW corner of Sec. 29, N pine (probably white pine) (20", 24"); east between Sections 29 and 32 from SW corner of Sec. 29, 24.00 ch. leave "timber & enter small prairie;" 40.00 ch. bur oak (6", 8"); comments include "Sand 3rd rate" and timber noted was "R & N pine [red and white] & tamarac."

STAND 11. Location: Mahnomen Co., 8 $\frac{1}{2}$ mi. E of Mahnomen and $\frac{1}{4}$ mi. S; NW $\frac{1}{4}$ NW $\frac{1}{4}$ Sec. 5, T 144 N, R 40 W. Topography: Middle slope above small lake, E 20° aspect, 6% slope; moist sandy loam soil. Vegetation: Wet to mesic prairie with Andropogon gerardi (up to 4' tall), A. scoparius, Amorpha canescens, Petalostemum spp., and Solidago spp.; 2% bare ground. Remarks: Active gopher mounds being invaded by Symphoricarpos occidentalis; Populus tremuloides and Rhus glabra invading from upland; prairie unplowed but fall mowed annually. Survey records: September-October, 1874; comments include "rolling prairie," "first rate" soil. (See also Survey records, Stand 12.)

STAND 12. Location: Mahnomen Co., $8\frac{1}{2}$ mi. E of Mahnomen and $\frac{1}{2}$ mi. S; $SE\frac{1}{4}$ $SE\frac{1}{4}$ $NE\frac{1}{4}$ and $NE\frac{1}{4}$ $NE\frac{1}{4}$ $SE\frac{1}{4}$ Sec. 6, T 144 N, R 40 W. Topography: Hill above prairie and marsh land, E aspect, near top of hill, 3-20% slope; sandy loam soil. Vegetation: 60% canopy of Quercus macrocarpa (up to 20" X 50') with Tilia americana and Ulmus americana; well developed shrub layer; scattered forbs; no bare soil (litter covered). Remarks: Stand not disturbed for at least 25 years, and then only posts removed. Survey records: September-October, 1874; at SE corner of Sec. 6, oak (10", 14", 14", 20"), "timber oak & aspen;" at quarter corner between Sections 6 and 7, ash (10") and willow (4"), "timber elm, oak, & aspen;" S between Sections 5 and 6 from NE corner of Sec. 6, 44.00 ch. oak (6", 6"), timber "elm, oak, aspen, and sugar;" comments include "soil first rate" and "North $\frac{1}{2}$ rolling prairie" and "South $\frac{1}{2}$ timber."

STAND 13. Location: Norman Co., 4 mi. E of Syre and $1\frac{1}{8}$ mi. N; $NW\frac{1}{4}$ $SW\frac{1}{4}$ $SW\frac{1}{4}$ Sec. 17, T 143 N, R 43 W. Topography: Level land on top of moraine ridge; gravelly soil. Vegetation: 40% canopy of Quercus macrocarpa with Populus tremuloides and Acer negundo; well developed shrub understory; no bare soil. Remarks: Abandoned homesite with open grown oaks. Survey records: July, 1871; SW corner of Sec. 17 is in prairie; comments include "Timber oak and aspen;" "Aspen, hazel & willow undergrowth;" "high peak, highest of area."

STAND 14. Location: Clay Co., $3\frac{1}{2}$ mi. E of Glyndon and $1\frac{1}{2}$ mi. S; $SE\frac{1}{4}$ $NE\frac{1}{4}$ Sec. 16, T 139 N, R 46 W. Topography: Level land; sandy gravelly soil. Vegetation: Mesic prairie with Andropogon gerardi, A. scoparius and associated species; no bare soil. Remarks: Large boulders on surface; never been plowed, but fall mowed annually; drainage ditches within 200 yards; shrubs and trees (including Cornus stolonifera [one individual 13 years old], Salix spp. [4 individuals 12 to 49+ years old], and Populus tremuloides) becoming established around rocks and in low places. Survey records: September, 1870; "level prairie."

STAND 15. Location: Polk Co., 1 mi. S of Fertile and $\frac{1}{2}$ mi. W; Agassiz Dunes Nature Conservancy Area; $W\frac{1}{2}$ $NE\frac{1}{4}$ Sec. 32, T 147 N, R 44 W. Topography; Sand dunes with all aspects, level to 40% slope, some old blowouts; very fine sand. Vegetation: Primarily Quercus macrocarpa (4-8" X 30') savanna with prairie understory and Populus tremuloides invading low areas; Juniperus horizontalis common; 5% bare soil. Remarks: The area has been grazed in the past, but not recently; very few active blowouts. Survey records: August-September, 1872; quarter corner between Sections 29 and 32, oak (10", 10"); NW corner of Sec. 32, oak (5", 6", 6", 8"); quarter corner between Sections 32 and 33, "Brush prairie;" comments on north side of Sec. 32, "Surface hilly and broken;" "Soil 2nd rate;" "Scattering oaks, patches of prairie & brush."

STAND 16. Location: Marshall Co., 3 mi. S of Karlstad;
 NE $\frac{1}{4}$ SE $\frac{1}{4}$ Sec. 5, T 158 N, R 45 W. Topography: Low, level,
 standing water in places during wet seasons: gravelly soil.
 Vegetation: 45% canopy of Populus tremuloides (4-7" X 55')
 with some Quercus macrocarpa (4-5" X 25'); dense shrub under-
 story (4-7' tall); no bare soil. Remarks: Railroad immediately
 W of stand may influence water standing in wet seasons; not
 disturbed recently, but evidently in past (cause undetermined).
 Survey records: February-March, 1877; SE corner of Sec. 5,
 aspen (4", 5", 6", 7"); quarter corner between Sections 4 and 5,
 aspen (5", 6"); comments include "Land 2nd rate;" "Brush prairie &
 scattering aspen;" "Timber small aspen."

STAND 17. Location: Kittson Co., 5 mi. NE of Karlstad;
 NE $\frac{1}{4}$ SW $\frac{1}{4}$ Sec. 3, T 159 N, R 45 W. Topography: Flat land above
 marshes; sandy soil. Vegetation: 30% canopy of Quercus
macrocarpa, with some small Populus tremuloides; shrub cover
 30-40%; prairie ground cover; no bare soil. Remarks: Evidently
 this stand was cut over as evidenced by the Q. macrocarpa
 sprouts, but several mature oaks left; P. tremuloides also
 removed. Survey records: April, 1877; SW corner of Sec. 3
 is "marsh land;" E from SW corner of Sec. 3, 20.00 ch. leave
 "open marsh enter tamarac;" 25.00 ch. leave "tamarac & enter
 oak;" 34.40 ch. aspen (10", 12"); quarter corner between Sections
 3 and 4 is "marsh;" "upland 2nd rate [soil]." (Evidently the
 tamarack is between marsh and oak/aspens upland in this section).

5.2 Additional Vegetation Descriptions

5.21. Tree Reproduction and Shrub Data

Stand 2, Bur oak woods. Density of reproduction (trees 1" or less DBH) and shrubs by plot size and per m²; density and height range by two age classes.

Species	Number					Number per 8m ² plot and range of height by age classes			
	Per plot				Per m ² *				
	Plot size, m ²					4 years or less		Over 4 years	
	1	2	4	8		No.	Ht.	No.	Ht.
<i>Fraxinus pennsylvanica</i>	0	0	2	3	0.0938	3	10-20"	0	-
<i>Ostrya virginiana</i>	0	0	2	5	0.1562	2	4-20"	3	1.5-6'
<i>Prunus virginiana</i>	2	3	12	16	0.5000	11	4-36"	5	1-4'
<i>Quercus macrocarpa</i>	4	4	12	19	0.5938	16	4-10"	3	8"
<i>Ribes cynosbati</i>	4	14	29	44	1.3750	41	4-25"	3	3'
<i>Symphoricarpos occidentalis</i>	13	18	30	42	1.3215	28	6-36"	14	1-3'
<i>Tilia americana</i>	4	4	9	17	0.5312	8	4-6"	9	1-4'
<i>Ulmus americana</i>	1	2	8	10	0.3125	10	3-24"	0	-
<i>Xanthoxylum americanum</i>	9	14	24	39	1.2188	14	4-20"	25	1.5-7'

* Based on 8m² plot size.

Basis: 4 plots of each size given.

Stand 5, Bur oak woods. Density of reproduction (trees 1" or less DBH) and shrubs by plot size and per m²; density and height range by two age classes.

Species	<div>Number</div>					Number per 8m ² plot and range of height by age classes			
	Per plot				Per m ² *	4 years or less		Over 4 years	
	Plot size, m ²					No.	Ht.	No.	Ht.
	1	2	4	8					
Acer negundo	0	0	0	1	0.0312	1	24"	0	-
Fraxinus pennsylvanica	0	3	3	3	0.0938	3	18"	0	-
Prunus virginiana	0	0	0	2	0.0625	1	18"	1	4'
Quercus macrocarpa	0	2	2	5	0.1562	5	4-18"	0	-
Ribes cynosbati	38	97	151	218	6.8125	211	3-36"	7	3-5'
Symphoricarpos occidentalis	29	58	94	153	4.7812	149	6-20"	4	1.5-2'
Ulmus americana	1	3	3	3	0.0938	3	10-15"	0	-
Xanthoxylum americanum	1	2	4	10	0.3125	9	10-30"	1	4'

* Based on 8m² plot size.

Basis: 4 plots of each size given.

Stand 6, Bur oak woods. Density of reproduction (trees 1" or less DBH) and shrubs by plot size and per m²; density and height range by two age classes.

Species	Number					Number per 8m ² plot and range of height by age classes			
	Per plot				Per m ² *	4 years or less		Over 4 years	
	Plot size, m ²					No.	Ht.	No.	Ht.
	1	2	4	8					
Acer negundo	4	11	12	15	0.4688	8	6-60"	7	2-8'
Fraxinus pennsylvanica	8	12	23	37	1.1562	37	6-12"	0	-
Prunus virginiana	1	1	6	11	0.3438	8	6-48"	3	5-6'
Quercus macrocarpa	0	0	1	4	0.1250	3	2"	1	2'
Ribes cynosbati	3	12	18	30	0.9375	28	3-24"	2	1-4'
Rubus occidentalis	9	16	27	39	1.2188	39	12-24"	0	-
Symphoricarpos occidentalis	6	8	12	13	0.4062	12	6-24"	1	3'
Xanthoxylum americanum	3	10	15	31	0.9688	29	6-36"	2	3-4'

* Based on 8m² plot size.

Basis: 4 plots of each size given.

Stand 7, Bur oak savanna. Density of reproduction (trees 1" or less DBH) and shrubs by plot size and per m²; density and height range by two age classes.

Species	<div>Number</div>					Number per 8m ² plot and range of height by age classes			
	Per plot				Per m ² *	4 years or less		Over 4 years	
	Plot size, m ²					No.	Ht.	No.	Ht.
	1	2	4	8					
Quercus macrocarpa	1	2	2	8	0.25	0	-	8	0.5-6'

* Based on 8m² plot size.

Basis: 4 plots of each size given.

Stand 10, Bur oak woods. Density of reproduction (trees 1" or less DBH) and shrubs by plot size and per m²; density and height range by two age classes.

Species	Number					Number per 8m ² plot and range of height by age classes			
	Per plot				Per m ² *	4 years or less		Over 4 years	
	Plot size, m ²					No.	Ht.	No.	Ht.
	1	2	4	8					
Amelanchier humilis	6	10	11	26	0.8125	15	6-48"	11	3-6'
Corylus americana	12	26	40	84	2.6250	79	10-48"	5	5'
Crataegus rotundifolia	1	1	2	5	0.1562	5	10-36"	0	-
Lonicera canadensis	7	10	11	13	0.4062	13	4-36"	0	-
Prunus virginiana	3	6	16	35	1.0938	30	6-36"	5	2-6'
Rhus radicans	0	3	13	30	0.9375	30	6-18"	0	-
Rosa blanda	4	8	11	18	0.5625	17	6-48"	1	5'
Rubus strigosus	0	0	0	2	0.0625	2	12-24"	0	-
Symphoricarpos occidentalis	9	15	30	71	2.2188	61	6-48"	10	3-4'

* Based on 8m² plot size.

Basis: 4 plots of each size given.

Stand 11, Prairie. Density of reproduction (trees 1" or less DBH) and shrubs by plot size and per m²; density and height range by two age classes.

Species	<div>Number</div>					Number per 8m ² plot and range of height by age classes			
	Per plot				Per m ² *	4 years or less		Over 4 years	
	Plot size, m ²					No.	Ht.	No.	Ht.
	1	2	4	8					
Rosa arkansana	3	4	4	6	0.1875	6	6"	0	-
Symphoricarpos occidentalis	1	1	9	9	0.2812	9	6"	0	-

* Based on 8m² plot size.

Basis: 4 plots of each size given.

Stand 12, Bur oak woods. Density of reproduction (trees 1" or less DBH) and shrubs by plot size and per m²; density and height range by two age classes.

Species	Number					Number per 8m ² plot and range of height by age classes			
	Per plot				Per m ² *	4 years or less		Over 4 years	
	Plot size, m ²					No.	Ht.	No.	Ht.
	1	2	4	8					
<i>Acer negundo</i>	11	29	50	95	2.9688	93	2-24"	2	2'
<i>Cornus racemosa</i>	1	4	6	6	0.1875	7	30-48"	1	6'
<i>Cornus stolonifera</i>	0	0	3	3	0.0938	3	4-12"	0	-
<i>Crataegus punctata</i>	0	0	6	14	0.4375	13	3-30"	1	3'
<i>Fraxinus pennsylvanica</i>	2	2	7	9	0.2812	8	2-24"	1	6'
<i>Populus tremuloides</i>	0	0	2	2	0.0625	2	4"	0	-
<i>Prunus virginiana</i>	2	2	7	13	0.4062	9	2-42"	4	2-7'
<i>Rhus radicans</i>	10	40	71	119	3.7188	112	3-24"	7	1.25-2.5'
<i>Rubus occidentalis</i>	2	2	4	4	0.1250	4	12"	0	-
<i>Sambucus pubens</i>	0	3	3	4	0.1250	4	3-6"	0	-
<i>Symphoricarpos occidentalis</i>	13	16	23	30	0.9375	30	6-36"	0	-

* Based on 8m² plot size.

Basis: 4 plots of each size given.

Stand 15, Bur oak savanna. Density of reproduction (trees 1" or less DBH) and shrubs by plot size and per m²; density and height range by two age classes.

Species	Number					Number per 8m ² plot and range of height by age classes			
	Per plot				Per m ² *				
	Plot size, m ²					4 years or less		Over 4 years	
	1	2	4	8		No.	Ht.	No.	Ht.
Prunus virginiana	0	0	0	4	0.1250	4	12-24"	0	-
Salix interior	4	9	12	12	0.3750	12	8-12"	0	-
Spiraea alba	0	2	6	16	0.5000	16	6-12"	0	-

* Based on 8m² plot size.

Basis: 4 plots of each size given.

Stand 16, Aspen-oak woods. Density of reproduction (trees 1" or less DBH) and shrubs by plot size and per m²; density and height range by two age classes.

Species	<div>Number</div>					Number per 8m ² plot and range of height by age classes			
	Per plot				Per m ² *	4 years or less		Over 4 years	
	Plot size, m ²					No.	Ht.	No.	Ht.
	1	2	4	8					
Amelanchier alnifolia	0	3	6	9	0.2812	6	6-72"	3	6'
Cornus alternifolia	1	4	5	6	0.1875	6	12-36"	0	-
Cornus racemosa	3	5	9	25	0.7812	19	12-60"	6	5-6'
Cornus stolonifera	6	8	18	25	0.7812	20	12-72"	5	4-8'
Corylus americana	26	55	88	152	4.7500	112	3-60"	40	3-7'
Populus balsamifera	0	1	2	3	0.0938	3	48-60"	0	-
Prunus virginiana	7	7	11	12	0.3750	10	6-48"	2	6-7'
Quercus macrocarpa	1	1	1	3	0.0938	3	6-24"	0	-
Rosa blanda	3	6	12	21	0.6562	19	12-48"	2	4-5'
Salix sp.	0	0	0	2	0.0625	0	-	2	3-7'
Symphoricarpos occidentalis	3	4	6	9	0.2812	8	4-60"	1	4'

* Based on 8m² plot size.

Basis: 4 plots of each size given.

5.22 Herb and Vine Data

Stand 2, Bur oak woods. Frequency (%) of herbs and vines by plot sizes.

Species	Plot size, m ²			
	1	2	4	8
<i>Amphicarpa bracteata</i>	25	25	25	25
<i>Aster</i> sp.	75	75	75	75
<i>Carex brevior</i>	50	50	50	50
<i>Celastrus scandens</i>	0	0	0	25
<i>Circaea quadrisulcata</i>	50	50	50	50
<i>Galium boreale</i>	25	25	25	25
<i>Galium triflorum</i>	50	50	50	50
<i>Hydrophyllum virginianum</i>	100	100	100	100
<i>Lathyrus venosus</i>	25	25	25	25
<i>Mentha arvensis</i>	25	25	25	25
<i>Osmorhiza claytoni</i>	25	25	25	25
<i>Parthenocissus quinquefolia</i>	0	50	50	50
<i>Poa pratensis</i>	75	75	75	100
<i>Polygonatum canaliculatum</i>	25	25	25	50
<i>Potentilla</i> sp.	25	25	25	25
<i>Smilacina stellata</i>	50	50	50	50
<i>Taraxacum officinale</i>	0	25	25	25
<i>Thalictrum dioicum</i>	25	25	25	25
<i>Viola</i> sp.	75	75	75	75
<i>Vitis riparia</i>	0	50	50	75

Basis: 4 plots of each size given.

Stand 5, Bur oak woods. Frequency (%) of herbs and vines by plot sizes.

Species	Plot size, m ²			
	1	2	4	8
Anemone canadensis	50	75	75	75
Bidens frondosa	0	0	25	50
Galium boreale	25	25	25	25
Galium triflorum	50	75	75	100
Geum sp.	0	0	0	50
Hackelia virginiana	0	0	0	50
Hydrophyllum virginianum	0	0	25	25
Leonurus cardiaca	0	0	0	25
Melilotus officinalis	25	25	25	25
Parthenocissus quinquefolia	0	0	25	25
Poa pratensis	50	75	75	75
Polygonatum canaliculatum	0	25	25	25
Viola sp.	0	0	25	25

Basis: 4 plots of each size given.

Stand 6, Bur oak woods. Frequency (%) of herbs and vines by plot sizes.

Species	Plot size, m ²			
	1	2	4	8
<i>Amphicarpa bracteata</i>	0	25	25	25
<i>Arctium minus</i>	25	25	25	25
<i>Arisaema atrorubens</i>	0	0	0	25
<i>Laportea canadensis</i>	50	50	50	50
<i>Parthenocissus quinquefolia</i>	75	75	75	100
<i>Phryma leptostachya</i>	0	25	25	25
<i>Poa pratensis</i>	25	25	25	50
<i>Smilacina stellata</i>	0	25	25	25
<i>Thalictrum dioicum</i>	0	0	25	25
<i>Vicia americana</i>	25	25	25	25
<i>Viola</i> sp.	0	0	0	25

Basis: 4 plots of each size given.

Stand 7, Bur oak savanna. Frequency (%) of herbs and vines by plot sizes.

Species	Plot size, m ²			
	1	2	4	8
<i>Ambrosia psilostachya</i>	75	75	75	75
<i>Amorpha canescens</i>	50	50	75	75
<i>Andropogon gerardi</i>	25	25	25	50
<i>Andropogon scoparius</i>	50	75	75	100
<i>Anemone patens</i>	0	0	25	25
<i>Aristida tuberculosa</i>	25	25	25	25
<i>Artemisia ludoviciana</i>	0	0	0	25
<i>Aster ericoides</i>	0	0	25	25
<i>Calamovilfa longifolia</i>	50	100	100	100
<i>Carex</i> sp.	100	100	100	100
<i>Heuchera richardsoni</i>	0	0	25	25
<i>Koeleria cristata</i>	0	25	25	25
<i>Liatris aspera</i>	0	0	0	25
<i>Lithospermum canescens</i>	0	25	25	25
<i>Lithospermum caroliniense</i>	25	25	75	75
<i>Panicum capillare</i>	50	50	50	100
<i>Panicum virgatum</i>	0	0	0	25
<i>Poa pratensis</i>	100	100	100	100
<i>Solidago nemoralis</i>	25	25	25	50
<i>Stipa comata</i>	0	0	25	25
<i>Tragopogon dubius</i>	25	25	25	25

Basis: 4 plots of each size given.

Stand 10, Bur oak woods. Frequency (%) of herbs and vines by plot sizes.

Species	Plot size, m ²			
	1	2	4	8
<i>Agastache foeniculum</i>	0	0	0	25
<i>Anemone riparia</i>	0	25	25	25
<i>Bromus inermis</i>	25	50	50	50
<i>Carex pensylvanica</i>	100	100	100	100
<i>Elymus canadensis</i>	0	75	75	75
<i>Fragaria virginiana</i>	25	50	50	50
<i>Galium boreale</i>	50	75	75	75
<i>Maianthemum canadense</i>	50	50	50	75
<i>Parthenocissus quinquefolia</i>	0	0	0	25
<i>Poa pratensis</i>	75	75	75	75
<i>Polygonatum canaliculatum</i>	25	25	25	25
<i>Prenanthes alba</i>	25	25	25	25
<i>Smilax herbacea</i>	0	25	50	50
<i>Smilacina stellata</i>	25	25	25	25
<i>Thalictrum dioicum</i>	50	50	50	50
<i>Vicia americana</i>	0	0	0	25
<i>Viola</i> sp.	0	0	25	25

Basis: 4 plots of each size given.

Stand 11, Prairie. Frequency (%) of herbs and vines by plot sizes.

Species	Plot size, m ²			
	1	2	4	8
<i>Allium stellatum</i>	100	100	100	100
<i>Amorpha canescens</i>	75	75	75	75
<i>Andropogon gerardi</i>	100	100	100	100
<i>Andropogon scoparius</i>	75	75	75	75
<i>Anemone cylindrica</i>	50	100	100	100
<i>Antennaria neglecta</i> var. <i>neglecta</i>	25	25	25	25
<i>Apocynum androsaemifolium</i>	25	25	50	50
<i>Aster puniceus</i>	75	75	75	100
<i>Astragalus crassicaupus</i>	100	100	100	100
<i>Bouteloua curtipendula</i>	0	25	25	25
<i>Campanula rotundifolia</i>	25	50	50	50
<i>Carex</i> sp.	75	100	100	100
<i>Elymus canadensis</i>	25	50	75	100
<i>Erigeron strigosus</i>	0	25	25	25
<i>Fragaria virginiana</i>	50	50	50	50
<i>Galium boreale</i>	75	75	75	100
<i>Heliopsis helianthoides</i>	25	25	25	25
<i>Koeleria cristata</i>	50	100	100	100
<i>Lathyrus venosus</i>	0	0	0	25
<i>Liatrus aspera</i>	50	100	100	100
<i>Lithospermum canescens</i>	50	50	50	50
<i>Melilotus alba</i>	25	25	25	25
<i>Monarda fistulosa</i>	25	25	25	25
<i>Panicum liebergii</i>	100	100	100	100
<i>Penstemon gracilis</i>	50	50	50	50
<i>Petalostemum candidum</i>	50	75	75	75
<i>Petalostemum purpureum</i>	75	75	100	100
<i>Phlox pilosa</i>	25	25	25	25
<i>Psoralea argophylla</i>	50	50	50	50
<i>Psoralea cuspidata</i>	25	25	25	25
<i>Ranunculus repens</i>	25	25	50	50
<i>Smilacina stellata</i>	0	50	50	50
<i>Solidago canadensis</i>	0	0	25	25
<i>Solidago rigida</i>	50	50	50	50
<i>Stipa spartea</i>	100	100	100	100
<i>Thalictrum dasycarpum</i>	50	75	75	75
<i>Veronicastrum virginicum</i>	25	25	50	50
<i>Zizia aptera</i>	25	50	50	50
<i>Zizia aurea</i>	0	0	75	75
<i>Zygadenus glaucus</i>	25	25	25	25

Basis: 4 plots of each size given.

Stand 12, Bur oak woods. Frequency (%) of herbs and vines by plot sizes.

Species	Plot size, m ²			
	1	2	4	8
<i>Amphicarpa bracteata</i>	50	50	50	75
<i>Apocynum androsaemifolium</i>	0	0	25	25
<i>Aralia nudicaulis</i>	50	50	50	50
<i>Carex</i> sp.	25	25	50	75
<i>Desmodium glutinosum</i>	50	75	75	100
<i>Elymus canadensis</i>	0	0	50	50
<i>Fragaria virginiana</i>	0	0	0	25
<i>Heracleum lanatum</i>	0	0	0	25
<i>Parthenocissus quinquefolia</i>	25	25	25	75
<i>Phryma leptostachya</i>	0	0	25	25
<i>Sanguinaria canadensis</i>	0	50	50	75
<i>Smilacina racemosa</i>	0	0	0	50
<i>Smilacina stellata</i>	0	0	0	25
<i>Thalictrum dioicum</i>	50	75	75	75
<i>Trillium grandiflorum</i>	0	25	25	25
<i>Uvularia grandiflora</i>	100	100	100	100
<i>Vicia americana</i>	25	25	25	25
<i>Viola</i> sp.	25	50	50	50
<i>Zizia aurea</i>	25	75	75	75

Basis: 4 plots of each size given.

Stand 14, Prairie. Frequency (%) of herbs and vines by plot sizes.

Species	Plot size, m ²			
	1	2	4	8
<i>Agropyron smithii</i>	25	50	50	75
<i>Agrostis stolonifera</i>	100	100	100	100
<i>Andropogon gerardi</i>	100	100	100	100
<i>Andropogon scoparius</i>	75	75	75	75
<i>Apocynum sibiricum</i>	0	0	25	25
<i>Asclepias ovalifolia</i>	0	0	25	25
<i>Aster ericoides</i>	0	0	25	25
<i>Aster ptarmicoides</i>	0	0	0	25
<i>Calamagrostis canadensis</i>	100	100	100	100
<i>Carex interior</i>	75	100	100	100
<i>Cirsium flodmanii</i>	0	0	0	50
<i>Equisetum kansanum</i>	0	0	0	25
<i>Liatris ligulistylis</i>	25	25	75	100
<i>Koeleria cristata</i>	25	50	50	75
<i>Penstemon gracilis</i>	0	0	25	25
<i>Petalostemum purpureum</i>	0	25	75	75
<i>Rudbeckia hirta</i>	25	50	75	75
<i>Solidago missouriensis</i>	25	50	100	100
<i>Taraxacum officinale</i>	0	25	25	25
<i>Thalictrum dasycarpum</i>	25	25	50	50
<i>Zizia aptera</i>	0	25	25	50
<i>Zizia aurea</i>	100	100	100	100

Basis: 4 plots of each size given.

Stand 15, Bur oak savanna. Frequency (%) of herbs and vines by plot sizes.

Species	Plot size, m ²			
	1	2	4	8
<i>Ambrosia psilostachya</i>	25	25	50	50
<i>Amorpha canescens</i>	75	75	75	75
<i>Andropogon gerardi</i>	0	25	50	50
<i>Andropogon scoparius</i>	25	25	25	50
<i>Anemone cylindrica</i>	25	25	25	25
<i>Artemisia ludoviciana</i>	50	50	50	50
<i>Aster</i> sp.	25	25	25	25
<i>Calamovilfa longifolia</i>	50	50	50	50
<i>Carex siccata</i>	0	0	25	25
<i>Koeleria cristata</i>	25	25	25	50
<i>Liatris aspera</i>	25	25	25	25
<i>Poa pratensis</i>	100	100	100	100
<i>Solidago missouriensis</i>	50	50	75	75
<i>Sorghastrum nutans</i>	25	25	25	25
<i>Stipa spartea</i>	25	25	25	25

Basis: 4 plots of each size given.

Stand 16, Aspen-oak woods. Frequency (%) of herbs and vines by plot sizes.

Species	Plot size, m ²			
	1	2	4	8
<i>Aster macrophyllus</i>	50	50	50	50
<i>Bromus inermis</i>	25	25	25	25
<i>Elymus canadensis</i>	75	75	75	75
<i>Fragaria virginiana</i>	50	50	75	75
<i>Ipomea purpurea</i>	25	25	25	50
<i>Maianthemum canadense</i>	75	75	100	100
<i>Sanicula marilandica</i>	50	100	100	100
<i>Smilacina stellata</i>	25	25	25	50
<i>Solidago rigida</i>	50	75	100	100
<i>Thalictrum dioicum</i>	25	75	75	75
<i>Veronicastrum virginicum</i>	0	0	0	25
<i>Zizia aurea</i>	75	75	75	75

Basis: 4 plots of each size given.

5.3 Soils Data

5.31 Soil Descriptions

Stand 2

General Site Characteristics:

Location - Jackson County, Minnesota, 7 mi. N and 4 mi. W of Jackson, Kilen Woods State Park; $W\frac{1}{2}$ $NE\frac{1}{4}$ Sec. 17 and $N\frac{1}{4}$ $SW\frac{1}{4}$ Sec. 16, T 103 N, R 35 W, 5th P.M.

Elevation - About 1425 feet.

Topography - Mostly level land intersected by large draw with perennial stream; N aspect, 0-5% slope, upland above draw.

Drainage - Well drained; moderate permeability.

Parent material - Calcareous loam till of Wisconsin age.

Vegetation - Bur oak woods (description found in Sections 2.221, 2.322, 5.1, and 5.2).

Erosion - None.

Climate - Mean annual temperature, 46°F; mean annual precipitation, 26 inches.

Soil family - Typic Hapludolls, fine-loamy, mixed, mesic; resembling the Clarion series.

Profile Description:

A soil pit was not dug at this stand as it was within a small state park. See Section 5.32 for a chemical and physical description of the soil.

Stand 5

General Site Characteristics:

Location - Lyon County, Minnesota, 14 mi. S of Marshall;
NW $\frac{1}{4}$ SW $\frac{1}{4}$ Sec. 11, T 109 N, R 40 W, 5th P.M.

Elevation - About 1450 feet.

Topography - Rolling with draws and small seasonal streams;
uplands above Cottonwood River; S 60° E aspect, 5-10% slope,
upper slope.

Drainage - Well drained; moderate permeability.

Parent material - Calcareous till of late Wisconsin age.

Vegetation - Bur oak woods (description found in Sections
2.221, 2.322, 5.1, and 5.2).

Erosion - None.

Climate - Mean annual temperature, 45°F; mean annual
precipitation, 25 inches.

Soil family - Udic Argiborolls, fine-loamy, mixed;
similar to the Forman series.

Profile Description:

01 1 - 0.5 inches. Litter; boundary abrupt-smooth.

02 0.5 - 1 inches. Decomposed litter; roots plentiful;
boundary abrupt-smooth.

A1 0 - 7 inches. Very dark brown (10 YR 2/2, moist) clay
loam (32 $\frac{1}{2}$ % sand, 35% silt, 32 $\frac{1}{2}$ % clay); dark gray (10 YR 4/1.5)
when dry; weak, medium, crumb structure; pores many; roots

plentiful; pH slightly acid (6.3); bulk density 1.14 g/cc;
7.28% organic matter; 78% base saturation; boundary abrupt-wavy.

A3 7 - 10 inches. Dark brown (10 YR 3/3, moist) clay loam
(36 $\frac{1}{2}$ % sand, 26 $\frac{1}{2}$ % silt, 37% clay); dark brown (10 YR 3/3) when
dry; weak, medium, crumb structure; pores many; roots few;
pH strongly acid (5.5); 5.17% organic matter; 76.1% base satu-
ration; boundary abrupt-wavy.

B1 10 - 12 inches. Dark brown (10 YR 3/3, moist) clay loam;
dark brown (10 YR 3/3) when dry; moderate, medium, crumb struc-
ture; pores common; roots few; boundary clear-wavy.

B2 12 - 18 inches. Dark brown (10 YR 3/3, moist) clay loam
(38 $\frac{1}{2}$ % sand, 23 $\frac{1}{2}$ % silt, 38% clay); yellowish brown (10 YR 5/4)
when dry; strong, medium, subangular blocky structure; pores
few; roots few; pH medium acid (5.8); 4.48% organic matter;
77.3% base saturation; boundary clear-wavy.

B3 18 - 20 inches. Dark brown (10 YR 4/3, moist) clay loam;
pale brown (10 YR 6/3) when dry; strong, medium, subangular blocky
structure; pores few; roots very few; boundary clear-wavy.

C1 20 - 27 inches. Yellowish brown (10 YR 5/5, moist) sandy
clay loam (45 $\frac{1}{2}$ % sand, 23 $\frac{1}{2}$ % silt, 31% clay); very pale brown
(10 YR 7/3) when dry; pores few; roots very few; pH neutral (7.3);
100% base saturation; boundary gradual-wavy.

C2 27 + inches. Yellowish brown (10 YR 5/6, moist) clay loam
(31 $\frac{1}{2}$ % sand, 34 $\frac{1}{2}$ % silt, 34% clay); very pale brown (10 YR 7/3.5)
when dry; pores few; roots very few; pH mildly alkaline (7.8);
100% base saturation.

Additional information on nutrient status and physical properties may be found in Section 5.32 . There is a gradation of organic matter on the ped faces from B1-C which is very pronounced. Rocks 1-2 inches in diameter are present in increasing amounts from B1-C.

Stand 6

General Site Characteristics:

Location - Pope County, Minnesota, 3 mi. NW of Glenwood;
NE $\frac{1}{4}$ SE $\frac{1}{4}$ Sec. 28, T 126 N, R 38 W, 5th P.M.

Elevation - About 1450 feet.

Topography - Top of rolling hills, above marshland and lakes; N 70° E aspect, 0-5% slope.

Drainage - Well drained; permeability moderate.

Parent material - Calcareous till of late Wisconsin age.

Vegetation - Bur oak woods (description found in Sections 2.221, 2.322, 5.1, and 5.2).

Erosion - None.

Climate - Mean annual temperature, 44°F; mean annual precipitation, 24 inches.

Soil family - Mollic Eutroboralfs, fine-loamy, mixed; resembling the Waukon series.

Profile Description:

- 01 1.5 - 0.5 inches. Leaf litter; boundary abrupt-smooth.
- 02 0.5 - 0 inches. Decomposed litter; boundary abrupt-smooth.
- A1 0 - 7 inches. Black (10 YR 2/1, moist) loam (36% sand, 39 $\frac{1}{2}$ % silt, 24 $\frac{1}{2}$ % clay); very dark gray (10 YR 3/1) when dry; moderate, fine, crumb structure; pores common; roots plentiful; pH slightly acid (6.2); bulk density 0.83 g/cc; 11.01% organic matter; 74.9% base saturation; boundary abrupt-wavy.

A3 7 - 8.5 inches. Very dark brown (10 YR 2/2, moist) clay loam (43% sand, 23½% silt, 33½% clay); very dark gray (10 YR 3/1) when dry; moderate, fine, crumb structure; pores common; roots plentiful; pH slightly acid (6.3); 5.91% organic matter; 76.4% base saturation; boundary abrupt-wavy.

B1 8.5 - 9.5 inches. Dark brown (10 YR 3/3, moist) clay loam; dark gray (10 YR 4/1) when dry; moderate, fine, angular blocky structure; pores common; roots few; boundary abrupt-wavy.

B2 9.5 - 18 inches. Dark brown (10 YR 4/3, moist) sandy clay loam (47% sand, 20% silt, 33% clay); dark gray brown (10 YR 4/2) when dry; moderate, fine, angular blocky structure; pores few; roots few; pH neutral (6.6); bulk density 1.21 g/cc; 3.63% organic matter; 81.5% base saturation; few limestone rocks 1-4 inches diameter; abrupt-irregular.

C 18 + inches. Dark yellowish brown (10 YR 4/4, moist) sandy clay loam (48% sand, 20½% silt, 31½% clay); yellowish brown (10 YR 5/4) when dry; moderate, fine, angular blocky structure; pH neutral (7.3); 98.8% base saturation; numerous limestone rocks 1-4 inches diameter.

Additional information on nutrient status and physical properties may be found in Section 5.32. At the 36 inch depth, the texture was clay (23% sand, 28% silt 49% clay), pH mildly alkaline (7.6), 99.0% base saturation.

Stand 7

General Site Characteristics:

Location - Anoka County, Minnesota, East Bethel, Allison Savanna Nature Conservancy Area; NE $\frac{1}{4}$ NE $\frac{1}{4}$ Sec. 2, T 133 N, R 23 W, 5th P.M.

Elevation - About 910 feet.

Topography - Rolling sand dunes; S aspect, 0-5% slope, upper slope.

Drainage - Excessively drained; rapid permeability.

Parent material - Aeolian fine sand.

Vegetation - Bur oak savanna (description found in Sections 2.221, 2.322, 5.1, and 5.2).

Erosion - None or some dune blowouts.

Climate - Mean annual temperature, 43°F; mean annual precipitation, 30 inches.

Soil family - Typic Udipsamments, mixed, frigid; resembling the Sartell series.

Profile Description:

No soil pit was dug in this stand, as it is a small, intensively used area. See Section 5.32 for a chemical and physical description of the soil.

Stand 10

General Site Characteristics:

Location - Hubbard County, Minnesota, 3/4 mi. S of Hubbard;
W $\frac{1}{2}$ SE $\frac{1}{4}$ SW $\frac{1}{4}$ Sec. 29, T 139 N, R 34 W, 5th P.M.

Elevation - About 1410 feet.

Topography - Level land above slightly rolling topography.

Drainage - Well drained; rapid permeability.

Parent material - Coarse sandy outwash.

Vegetation - Bur oak woods (description found in Sections
2.221, 2.322, 5.1, and 5.2).

Erosion - None.

Climate - Mean annual temperature, 40°F; mean annual
precipitation, 26 inches.

Soil family - Udic Haploborolls, sandy, mixed; resembles
the Hubbard series.

Profile Description;

01 3.5 - 2 inches. Leaf litter; boundary abrupt-wavy.

02 2 - 0 inches. Decomposed litter; roots plentiful; boundary
abrupt-smooth.

A1 0 - 3 inches. Black (7.5 YR 2/0, moist) sand (82% sand,
10% silt, 8% clay); black (10 YR 2/1) when dry; single grain to
weak, coarse, crumb structure; loose (dry), very friable (moist),
nonsticky (wet) consistence; pores few; roots plentiful; pH
slightly acid (6.3); 4.55% organic matter; 63.7% base saturation;

Boundary abrupt-wavy.

B1 3 - 7 inches. Very dark gray brown (10 YR 3/2, moist) sand; dark brown (10 YR 3.5/3) when dry; single grain to weak, coarse, crumb structure; loose (dry), very friable (moist), nonsticky (wet) consistence; pores few; roots plentiful; bulk density 1.27 g/cc; boundary clear-wavy.

B2 7 - 11 inches. Very dark gray brown (10 YR 3/2, moist) sand (84½% sand, 11½% silt, 4% clay); dark brown (10 YR 4/3) when dry; single grain to weak, coarse, crumb structure; loose (dry), very friable (moist), slightly sticky (wet) consistence; pores few; roots few; pH medium acid (6.0); 2.44% organic matter; 55.5% base saturation; boundary clear-irregular.

C 11 + inches. Dark yellowish brown (10 YR 4.5/4, moist) loamy sand (77% sand, 11% silt, 12% clay); yellowish brown (10 YR 5.5/8) when dry; single grain structure; loose (dry); very friable (moist), slightly sticky (wet) consistence; roots few; pH medium acid (5.8); bulk density 1.62 g/cc; 2.45% organic matter; 56% base saturation.

Additional information on nutrient status and physical properties may be found in Section 5.32 . At the 24 inch depth, the texture was a loamy sand (78% sand, 10% silt, 12% clay), with a pH medium acid (5.8), 1.87% organic matter, and 50.4% base saturation; at the 36 inch depth, the texture was a sand (91% sand, 3% silt, 6% clay), with a pH medium acid (6.0), 0.87% organic matter, and 42.7% base saturation.

Stand 11

General Site Characteristics:

Location - Mahnomen County, Minnesota, $8\frac{1}{2}$ mi. E of Mahnomen and $\frac{1}{4}$ mi. S; NW $\frac{1}{4}$ NW $\frac{1}{4}$ Sec. 5, T 144 N, R 40 W, 5th P.M.

Elevation - About 1320 feet.

Topography - Middle slope above small lake, E 20° S aspect, 6% slope.

Drainage - Moderately well drained; moderate permeability.

Parent material - Calcareous till of late Wisconsin age.

Vegetation - Prairie (description found in Sections 2.221, 2.322, 5.1, and 5.2).

Erosion - None.

Climate - Mean annual temperature, 40°F ; mean annual precipitation, 24 inches.

Soil family - Aquic Argiborolls, fine, montmorillonitic; resembles the Gonvick series.

Profile Description:

01 0.5 - 0.1 inches. Litter; boundary very abrupt-smooth.

02 0.1 - 0 inches. Decomposed litter, boundary very abrupt-smooth.

A1 0 - 12 inches. Black (7.5 YR 2/0, moist) clay loam (37% sand, 35% silt, 28% clay); black (10 YR 2/1) when dry; moderate, medium, crumb structure; weakly coherent (dry); friable (moist), slightly sticky (wet) consistence; pores common; roots plentiful;

pH slightly acid (6.5); bulk density 0.93 g/cc; 11.22% organic matter; 82.3% base saturation; boundary abrupt-wavy.

B1 12 - 16 inches. Very dark gray (10 YR 3/1.5, moist) clay (18% sand, 34 $\frac{1}{2}$ % silt, 47 $\frac{1}{2}$ % clay); dark gray (10 YR 4.5/1) when dry; moderate, medium crumb structure; slightly hard (dry), friable (moist), slightly sticky (wet) consistence; pH strongly acid (5.5); 7.86% organic matter; 100% base saturation; boundary abrupt-wavy.

B2 16 - 26 inches. Dark yellowish brown (10 YR 4.5/4, moist) silty clay loam (19% sand, 31% silt, 50% clay); light gray (10 YR 7/1.5) when dry; strong, coarse, subangular blocky structure; slightly hard (dry), firm (moist), sticky (wet) consistence; pH strongly acid (5.5); bulk density 1.34 g/cc; 100% base saturation; boundary abrupt-wavy.

B3 26 + inches. Yellowish brown (10 YR 5.2/4, moist) clay loam (24% sand, 33 $\frac{1}{2}$ % silt, 42 $\frac{1}{2}$ % clay); light gray (10 YR 7/2) when dry; moderate to strong; coarse, subangular blocky structure; slightly hard (dry), firm (moist), sticky (wet) consistence; pH strongly acid (5.3); 100% base saturation.

Additional information on nutrient status and physical properties may be found in Section 5.32.

Stand 12

General Site Characteristics:

Location - Mahnomen County, Minnesota, $8\frac{1}{2}$ mi. E of Mahnomen and $\frac{1}{4}$ mi. S; $SE\frac{1}{4}$ $SE\frac{1}{4}$ $NE\frac{1}{4}$ and $NE\frac{1}{4}$ $NE\frac{1}{4}$ $SE\frac{1}{4}$ Sec. 6, T 144 N, R 40 W, 5th P.M.

Elevation - About 1370 feet.

Topography - Hill above prairie and marsh land, E aspect, near top of hill, 3-20% slope.

Drainage - Well drained; moderate permeability.

Parent material - Calcareous till of late Wisconsin age.

Vegetation - Bur oak woods (description found in Sections 2.221, 2.322, 5.1, and 5.2).

Erosion - None.

Climate - Mean annual temperature, 40°F ; mean annual precipitation, 24 inches.

Soil family - Udic Argiborolls, fine, montmorillonitic.

Profile Description:

01 3 - 1 inches. Litter; boundary abrupt-smooth.

02 1 - 0 inches. Decomposed litter; roots plentiful; boundary abrupt-smooth.

A1 0 - 14 inches. Very dark brown (10 YR 2.5/2, moist) loam (39% sand, 34% silt, 27% clay); very dark gray (10 YR 3.5/1) when dry; weak, fine, crumb structure; friable (moist), slightly sticky (wet) consistence; pores common; roots plentiful; pH mildly

alkaline (7.4); bulk density 1.12 g/cc; 10.61% organic matter; 100% base saturation; boundary clear-wavy.

A3 14 - 23 inches. Very dark gray (10 YR 3/1, moist) clay (27% sand, 29% silt, 44% clay); brown (10 YR 5/3) when dry; moderate, medium, crumb texture; friable (moist), slightly sticky (wet) consistence; pores common; roots plentiful; pH neutral (7.1); bulk density 1.21 g/cc; 4.61% organic matter; 100% base saturation; boundary gradual-wavy.

B2 23 - 37 inches. Very dark gray brown (10 YR 3/2.5, moist) clay (26.5% sand, 33.5% silt, 40% clay); light gray (10 YR 7/2) when dry; moderate, medium, crumb structure; firm (moist), sticky (wet) consistence; pores few; roots few; pH mildly alkaline (7.6); 100% base saturation; boundary clear-wavy.

B3 37 + inches. Dark brown (10 YR 4/3, moist) clay (19% sand, 38% silt, 43% clay); yellowish brown (10 YR 5/4) when dry; moderate, coarse, subangular blocky structure; firm (moist), sticky (wet) consistence; pH mildly alkaline (7.7); 100% base saturation; some calcareous rocks present.

Additional information on nutrient status and physical properties may be found in Section 5.32.

Stand 13

General Site Characteristics:

Location - Norman County, Minnesota, 4 mi. E of Syre and 1 1/8 mi. N; NW $\frac{1}{4}$ SW $\frac{1}{4}$ SW $\frac{1}{4}$ Sec. 17, T 143 N, R 43 W, 5th P.M.

Elevation - About 1310 feet.

Topography - Level land on top of moraine ridge.

Drainage - Well drained; moderate permeability.

Parent material - Calcareous loamy till of late Wisconsin age.

Vegetation - Bur oak woods (description found in Sections 2.221, 2.322, 5.1, and 5.2).

Erosion - None to very little.

Climate - Mean annual temperature, 40°F; mean annual precipitation, 23 inches.

Soil family - Udic Haploborolls, fine-loamy, mixed; resembles the Barnes series.

Profile Description:

01 1 - 0.25 inches. Litter; boundary abrupt-smooth.

02 0.25 - 0 inches. Decomposed litter; boundary abrupt-smooth.

A1 0 - 7.5 inches. Black (10 YR 2/1, moist) sandy loam (52 $\frac{1}{2}$ % sand, 34 $\frac{1}{2}$ % silt, 13% clay); very dark gray (10 YR 3/1) when dry; moderate, medium, crumb structure; weakly coherent (dry), friable (moist), slightly sticky (wet) consistence; pores

common; roots plentiful; pH neutral (6.7); bulk density 1.11 g/cc; 9.50% organic matter; 77.2% base saturation; boundary clear-smooth.

B1 7.5 - 11 inches. Very dark brown (10 YR 2/2, moist) sandy loam (57% sand, 30% silt, 13 % clay); very dark gray brown (10 YR 3.5/2.5) when dry; moderate, medium, crumb structure, weakly coherent (dry), friable (moist), slightly sticky (wet) consistence; pores common; roots plentiful; pH slightly acid (6.2); 6.31% organic matter; 68.7% base saturation; boundary clear-wavy.

B2 11 - 22 inches. Dark brown (10 YR 3.5/3, moist) sandy loam (63% sand, 18% silt, 19% clay); dark yellowish brown (10 YR 4/4) when dry; moderate coarse, crumb to subangular blocky structure; slightly hard (dry), friable (moist), sticky (wet) consistence; pores common; roots plentiful; pH slightly acid (6.1); bulk density 1.32 g/cc; 2.31% organic matter; 68.9% base saturation; boundary clear-wavy.

B3 22 + inches. Brown (10 YR 5.5/3, moist) sandy clay loam (61% sand, 18% silt, 21% clay); light brownish gray (10 YR 6.5/2.5) when dry; moderate, medium, crumb structure; slightly hard (dry), friable (moist), sticky (wet) consistence; pores few; roots few; pH slightly acid (6.1); 70.9% base saturation.

Additional information on nutrient status and physical properties may be found in Section 5.32 . At the 36 inch depth, the texture was loam (45 $\frac{1}{2}$ % sand, 27 $\frac{1}{2}$ % silt, 27% clay) with a pH mildly alkaline (7.6) and a 100% base saturation.

Stand 14

General Site Characteristics:

Location - Clay County, Minnesota, $3\frac{1}{2}$ mi. E of Glyndon and $1\frac{1}{2}$ mi. S; SE $\frac{1}{4}$ NE $\frac{1}{4}$ Sec. 16, T 139 N, R 46 W, 5th P.M.

Elevation - About 940 feet.

Topography - Level land.

Drainage - Poorly to somewhat poorly drained; permeability moderate.

Parent material - Calcareous till of late Wisconsin age and lacustrine sediments.

Vegetation - Prairie (description found in Sections 2.221, 2.322, 5.1, and 5.2).

Erosion - None.

Climate - Mean annual temperature, 42°F; mean annual precipitation, 22 inches.

Soil family - Typic Haplaquolls, coarse-loamy, mixed, calcareous, frigid.

Profile Description:

01 1 - 0.5 inches. Litter; boundary very abrupt-smooth.

02 0.5 - 0 inches. Decomposed litter; boundary abrupt-smooth.

A1 0 - 8.5 inches. Black (2.5 Y 2/0, moist) sandy loam (63 $\frac{1}{2}$ % sand, 22% silt, 14 $\frac{1}{2}$ % clay); very dark gray (2.5 Y 3/0)

when dry; single grain structure; pores common; roots abundant;

pH moderately alkaline (7.9); bulk density 1.12 g/cc; 9.58%

organic matter; 100% base saturation; boundary abrupt-wavy.

A3 8.5 - 11 inches. Black (2.5 Y 2.5/0, moist) sandy loam (64% sand, 17½% silt, 18½% clay); gray (2.5 Y 5/0) when dry; single grain structure; pores common; roots plentiful; pH moderately alkaline (7.9); 6.11% organic matter; 100% base saturation; boundary clear-wavy.

B1 11 - 17 inches. Dark gray (10 YR 4/1, moist) sandy loam (69% sand, 12% silt, 19% clay); light gray (2.5 Y 6.5/0) when dry; single grain structure; pores few; roots plentiful; pH moderately alkaline (8.0); 3.79% organic matter; 100% base saturation; boundary clear-wavy.

B2 17 - 32.5 inches. Dark gray (10 YR 4/1, moist) loamy sand (77½% sand, 8½% silt, 14% clay); gray (2.5 Y 6/0) when dry; single grain structure; pores common; roots few; pH moderately alkaline (8.1); 2.16% organic matter; 100% base saturation; boundary abrupt-wavy.

C 32.5 + inches. Dark brown (10 YR 4/3, moist) sand (89% sand, 3½% silt, 7½% clay); light brownish gray (10 YR 6/2) when dry; single grain structure; pores few; roots few; pH moderately alkaline (8.2); 0.63% organic matter; 100% base saturation.

Additional information on nutrient status and physical properties may be found in Section 5.32. Boulders are scattered over the surface and throughout the profile. Following heavy rains water table is at 2-3 feet below surface.

Stand 15

General Site Characteristics:

Location - Polk County, Minnesota, 1 mi. S of Fertile and $\frac{1}{2}$ mi. W, Agassiz Dunes Nature Conservancy Area; $W\frac{1}{2}$ NE $\frac{1}{4}$ Sec. 32, T 147 N, R 44 W, 5th P.M.

Elevation - About 1120 feet.

Topography - Sand dunes with all aspects, level to 40% slope, soil examined on level land.

Drainage - Excessively drained; rapid permeability.

Parent material - Aeolian fine sand.

Vegetation - Bur oak savanna (description found in Sections 2.221, 2.332, 5.1, and 5.2).

Erosion - None at soil pit to very severely eroded in dune blowouts.

Climate - Mean annual temperature, 40°F; mean annual precipitation, 22 inches.

Soil family - Typic Udipsamments, mixed, frigid.

Profile Description:

01 0.5 - 0.25 inches. Litter; boundary very abrupt-smooth.

02 0.25 - 0 inches. Decomposed litter; boundary abrupt-smooth.

A1 0 - 3.5 inches. Dark yellowish brown (10 YR 3/4, moist) sand (94% sand, 3% silt, 3% clay); dark gray brown (10 YR 4/2) when dry; single grain structure; roots abundant; pH slightly

acid (6.2); 1.59% organic matter; 51.4% base saturation; boundary abrupt-wavy.

A3 3.5 - 6 inches. Dark yellowish brown (10 YR 4/4, moist) sand (95% sand, 1 $\frac{1}{2}$ % silt, 3 $\frac{1}{2}$ % clay); brown (10 YR 5/3) when dry; single grain structure; roots abundant; pH slightly acid (6.2); bulk density 1.57 g/cc; 1.03% organic matter; 40.6% base saturation; boundary clear-wavy.

B1 6 - 8.5 inches. Dark yellowish brown (10 YR 4/4, moist) sand (94 $\frac{1}{2}$ % sand, 2% silt, 3 $\frac{1}{2}$ % clay); brown (10 YR 5/3) when dry; single grain structure; roots abundant; pH slightly acid (6.2); 0.61% organic matter; 40.2% base saturation; boundary clear-wavy.

B2 8.5 - 33.5 inches. Dark yellowish brown (10 YR 4/4, moist) sand (96% sand, $\frac{1}{2}$ % silt, 3 $\frac{1}{2}$ % clay); pale brown (10 YR 6/3) when dry; single grain structure; roots plentiful to few; pH slightly acid (6.4); bulk density 1.61 g/cc; 0.47% organic matter; 38.5% base saturation; boundary clear-wavy.

B3 33.5 + inches. Dark brown (10 YR 3/3, moist) sand (96% sand, 1% silt, 3% clay); brown (10 YR 5/3) when dry; single grain structure; roots very few; pH slightly acid (6.2); 0.35% organic matter; 40.2 % base saturation.

Additional information on nutrient status and physical properties may be found in Section 5.32.

Stand 16

General Site Characteristics:

Location - Marshall County, Minnesota, 3 mi. S of Karlstad;
NE $\frac{1}{4}$ SE $\frac{1}{4}$ Sec. 5, T 158 N, R 45 W, 5th P.M.

Elevation - About 1050 feet.

Topography - Low, level.

Drainage - Imperfectly or somewhat poorly drained; moderately slow permeability.

Parent material - Calcareous loam glacial till with perhaps some lacustrine sediments.

Vegetation - Aspen woods (description found in Sections 2.221, 2.332, 5.1, and 5.2).

Erosion - None.

Climate - Mean annual temperature, 38⁰F; mean annual precipitation, 20 inches.

Soil family - Typic Haplaquolls, fine-loamy, mixed, frigid; resembles the Roliss series.

Profile Description:

01 2 - 1 inches. Litter; boundary abrupt-smooth.

02 1 - 0 inches. Decomposed litter; roots plentiful; boundary abrupt-smooth.

A1 0 - 5 inches. Very dark gray brown (10 YR 3/2, moist) sandy clay loam (57 $\frac{1}{2}$ % sand, 21 $\frac{1}{2}$ % silt, 21% clay); gray (10 YR 5/1) when dry; weak, fine, crumb structure; pores common; roots

plentiful; pH neutral (6.8); bulk density 1.26 g/cc; 3.48% organic matter; 80.8% base saturation; boundary abrupt-wavy.

A2 5 - 7 inches. Dark gray brown (10 YR 4/2, moist) sandy loam ($64\frac{1}{2}\%$ sand, $19\frac{1}{2}\%$ silt, 16% clay); gray brown (10 YR 5/2) when dry; weak, medium, crumb structure; pores common; roots few; pH mildly alkaline (7.4); 2.66% organic matter; 100% base saturation; small rocks present; boundary abrupt-wavy.

B1 7 - 10 inches. Gray brown (10 YR 5/2, moist) clay loam ($41\frac{1}{2}\%$ sand, $30\frac{1}{2}\%$ silt, 28% clay); light brownish gray (10 YR 6/2) when dry; weak, medium, crumb structure; pores common; roots plentiful; pH moderately alkaline (8.0); 100% base saturation; several large rocks present; boundary abrupt-wavy.

B2 10 + inches. Pale brown (10 YR 6/3, moist) clay loam ($41\frac{1}{2}\%$ sand, $30\frac{1}{2}\%$ silt, 28% clay); light gray (10 YR 7.5/1) when dry; weak, medium crumb structure; pores few; roots few; pH moderately alkaline (8.1); bulk density 1.45 g/cc; 100% base saturation; many rocks present.

Additional information on nutrient status and physical properties may be found in Section 5.32 . The soil at 24 inches and below was too rocky to dig the soil pit deeper; the texture at 24 inches was clay loam ($41\frac{1}{2}\%$ sand, $30\frac{1}{2}\%$ silt, and 28% clay).

5.32 Soil Physical and Chemical Properties

Stand 2, Bur oak woods. Soil physical and chemical properties.

Property	Soil Depth, inches				
	0-6	6-12	12-18	18-24	36
Bulk density (g/cc)	1.212	-	-	-	-
Texture					
% Sand	39	39	41	42	44.5
% Silt	37	31	29	26	24.5
% Clay	24	30	30	32	31
Moisture properties					
15 Bar moist. (%)	15.93	12.11	11.15	10.95	11.03
1/3 Bar moist. (%)	27.10	25.60	24.70	24.35	24.15
Avail. moist. (%)	11.17	13.49	13.55	13.40	13.12
Organic matter (%)	6.89	4.75	3.67	3.07	2.53
pH	6.2	5.6	5.6	5.5	5.3
Nitrogen (%)	0.219	0.132	0.091	0.063	0.042
Phosphorus (ppm)	9.5	4.1	2.8	2.0	3.9
Potassium (ppm)	81.3	43.5	46.3	44.5	45.7
Calcium (ppm)	3587	2633	2677	2193	2067
Magnesium (ppm)	445	429	457	481	493
Sodium (ppm)	15.5	21.3	24.1	26.4	30.1
Hydrogen (ppm)	80.9	87.1	74.7	68.6	63.8
Base saturation (%)	73.1	65.9	70.2	68.9	69.9
Bases (meq/100g)	21.87	16.90	17.37	15.23	14.64
CEC (meq/100g)	29.89	25.54	24.78	21.95	20.96

Basis: Average of 3 replications, except bulk density with 6 at 6 inches.

Stand 5, Bur oak woods. Soil physical and chemical properties.

Property	Soil Depth, inches				
	0-7	7-12	12-18	18-24	36
Bulk density (g/cc)	1.190	-	1.319	-	-
Texture					
% Sand	34	36	36.5	38.5	33
% Silt	34	30	27.5	29	32.5
% Clay	32	34	36	32.5	34.5
Moisture properties					
15 Bar moist. (%)	19.82	16.45	14.96	13.07	11.98
1/3 Bar moist. (%)	29.10	26.74	26.66	25.40	27.03
Avail. moist. (%)	9.28	10.29	11.70	12.33	19.05
Organic matter (%)	7.79	5.52	4.62	4.39	4.22
pH	6.4	6.1	6.3	7.2	7.7
Nitrogen (%)	0.237	0.141	0.105	0.081	0.048
Phosphorus (ppm)	10.2	6.1	4.9	2.6	3.4
Potassium (ppm)	87.7	61.5	53.8	47.9	36.1
Calcium (ppm)	4078	3268	3207	7687	9667
Magnesium (ppm)	558	476	447	409	388
Sodium (ppm)	31.1	32.1	29.9	28.0	33.0
Hydrogen (ppm)	82.3	74.0	63.0	0.0	0.0
Base saturation (%)	75.7	73.3	76.0	100.0	100.0
Bases (meq/100g)	25.34	20.55	19.98	42.05	51.76
CEC (meq/100g)	33.50	27.90	26.23	42.05	51.76

Basis: Average of 3 replications, except bulk density with 6 at 6 inches and 4 at 18 inches.

Stand 6, Bur oak woods. Soil physical and chemical properties.

Property	Soil Depth, inches				
	0-6	6-12	12-18	18-24	36
Bulk density (g/cc)	0.833	-	1.211	-	-
Texture					
% Sand	33.5	36	38.5	43.5	31
% Silt	44	29.5	25.5	24	29
% Clay	22.5	34.5	36	32.5	40
Moisture properties					
15 Bar moist. (%)	27.83	19.30	16.88	15.40	14.38
1/3 Bar moist. (%)	36.36	28.70	27.70	26.43	28.84
Avail. moist. (%)	8.53	9.40	10.82	11.03	14.46
Organic matter (%)	12.23	6.66	4.83	4.00	3.19
pH	6.0	5.9	6.0	6.9	7.5
Nitrogen (%)	0.416	0.196	0.121	0.093	0.063
Phosphorus (ppm)	26.9	11.2	6.8	3.0	1.6
Potassium (ppm)	103.7	74.6	65.3	62.7	45.7
Calcium (ppm)	5713	3533	2913	6933	9747
Magnesium (ppm)	745	676	696	644	716
Sodium (ppm)	16.9	24.1	29.5	33.0	32.5
Hydrogen (ppm)	147.4	95.3	76.1	27.4	4.1
Base saturation (%)	70.8	71.4	72.8	90.9	99.2
Bases (meq/100g)	38.71	25.97	20.59	40.27	54.88
CEC (meq/100g)	49.65	32.97	28.13	42.99	55.29

Basis: Average of 3 replications, except bulk density with 6 at 6 inches and 2 at 18 inches.

Stand 7, Bur oak savanna. Soil physical and chemical properties.

Property	Soil Depth, inches					
	0-6	6-12	12-18	18-24	36	48
Bulk density (g/cc)	1.496	-	1.566	-	-	-
Texture						
% Sand	92.5	92	93	94	94.5	94.5
% Silt	2.5	2	2	0.5	0.5	0.5
% Clay	5	6	5	5.5	5	5
Moisture properties						
15 Bar moist. (%)	2.72	2.56	2.00	1.86	1.41	1.27
1/3 Bar moist. (%)	4.01	3.50	2.82	2.63	1.90	1.83
Avail. moist. (%)	1.29	0.94	0.82	0.77	0.49	0.56
Organic matter (%)	1.63	1.42	0.99	0.92	0.59	0.47
pH	5.2	5.3	5.5	5.5	5.6	5.9
Nitrogen (%)	0.057	0.043	0.028	0.024	0.008	0.004
Phosphorus (ppm)	14.1	11.5	13.6	13.1	10.0	7.3
Potassium (ppm)	5.9	5.4	3.5	3.5	3.1	3.6
Calcium (ppm)	161	144	116	109	143	183
Magnesium (ppm)	19.4	17.5	12.6	12.3	15.9	19.7
Sodium (ppm)	0.3	3.2	1.5	1.7	2.4	3.3
Hydrogen (ppm)	56.9	43.9	37.7	33.6	26.7	24.0
Base saturation (%)	14.8	18.3	16.7	17.2	25.1	34.7
Bases (meq/100g)	0.98	0.89	0.70	0.66	0.86	1.10
CEC (meq/100g)	6.63	5.25	4.44	3.99	3.51	3.48

Basis: Average of 3 replications, except bulk density with 6 at 6 inches and 2 at 18 inches.

Stand 10, Bur oak woods. Soil physical and chemical properties.

Property	Soil Depth, inches				
	0-6	6-12	12-18	18-24	36
Bulk density (g/cc)	1.371	-	1.618	-	-
Texture					
% Sand	82.5	84	80	80.5	91.5
% Silt	9	8.5	9	8.5	2.5
% Clay	8.5	7.5	11	11	6
Moisture properties					
15 Bar moist. (%)	7.34	4.34	4.26	3.75	1.82
1/3 Bar moist. (%)	9.60	7.17	7.91	7.18	2.74
Avail. moist. (%)	2.26	2.83	3.65	3.43	0.92
Organic matter (%)	3.70	2.43	2.08	1.72	0.90
pH	5.8	5.7	5.6	5.7	5.9
Nitrogen (%)	0.110	0.068	0.048	0.037	0.014
Phosphorus (ppm)	49.5	62.6	55.7	46.4	20.3
Potassium (ppm)	23.5	9.3	10.1	8.1	3.4
Calcium (ppm)	1377	978	1075	821	340
Magnesium (ppm)	138	125	100	84	34
Sodium (ppm)	9.5	6.2	5.3	7.4	2.6
Hydrogen (ppm)	65.1	52.8	59.6	50.1	29.5
Base saturation (%)	56.7	53.3	51.0	48.4	40.8
Bases (meq/100g)	8.12	5.97	6.25	4.85	2.00
CEC (meq/100g)	14.58	11.21	12.16	9.81	4.93

Basis: Average of 3 replications, except bulk density with 6 at 6 inches and 4 at 18 inches.

Stand 11, Prairie. Soil physical and chemical properties.

Property	Soil Depth, inches				
	0-6	6-12	12-18	18-24	36
Bulk density (g/cc)	0.993	-	1.335	-	-
Texture					
% Sand	38	33.5	25.5	29	24
% Silt	44	35.5	33	30	32
% Clay	18	31	41.5	41	44
Moisture properties					
15 Bar moist. (%)	25.70	18.42	13.79	11.80	13.13
1/3 Bar moist. (%)	36.42	31.35	28.55	25.77	24.67
Avail. moist. (%)	10.72	12.93	14.76	13.97	11.54
Organic matter (%)	12.50	8.58	7.52	6.21	5.99
pH	7.1	7.4	7.7	7.9	8.0
Nitrogen (%)	0.433	0.188	0.092	0.048	0.066
Phosphorus (ppm)	6.9	2.4	1.0	0.5	1.0
Potassium (ppm)	69.7	59.1	48.9	39.3	40.4
Calcium (ppm)	6813	7367	10027	9707	9787
Magnesium (ppm)	921	791	804	679	923
Sodium (ppm)	15.9	17.3	19.5	23.7	24.1
Hydrogen (ppm)	24.0	0.0	0.0	0.0	0.0
Base saturation (%)	94.1	100	100	100	100
Bases (meq/100g)	41.89	43.56	56.96	54.32	56.73
CEC (meq/100g)	44.27	43.56	56.96	54.32	56.73

Basis: Average of 3 replications, except bulk density with 8 at 6 inches and 4 at 18 inches.

Stand 12, Bur oak woods. Soil physical and chemical properties.

Property	Soil Depth, inches				
	0-6	6-12	12-18	18-24	36
Bulk density (g/cc)	1.024	-	1.209	-	-
Texture					
% Sand	41	36.5	31.5	25	26
% Silt	34	28.5	28.5	36	34.5
% Clay	25	35	40	39	39.5
Moisture properties					
15 Bar moist. (%)	16.40	12.28	14.48	14.33	13.86
1/3 Bar moist. (%)	26.41	23.78	27.93	37.42	26.62
Avail. moist. (%)	10.01	11.50	13.45	13.09	12.76
Organic matter (%)	8.11	5.37	4.81	4.23	4.18
pH	6.9	7.0	7.3	7.7	7.8
Nitrogen (%)	0.068	0.054	0.174	0.182	0.090
Phosphorus (ppm)	54.4	18.9	13.3	6.7	1.6
Potassium (ppm)	76.5	57.8	67.8	57.3	52.6
Calcium (ppm)	4675	6310	6795	8900	9195
Magnesium (ppm)	658	540	980	618	607
Sodium (ppm)	15.0	14.6	16.6	17.0	17.2
Hydrogen (ppm)	35.0	33.9	15.4	0.0	0.0
Base saturation (%)	90.5	88.5	95.4	100	100
Bases (meq/100g)	29.04	36.21	40.27	49.80	51.18
CEC (meq/100g)	32.52	39.57	41.80	49.80	51.18

Basis: Average of 3 replications, except bulk density with 6 at 6 inches and 3 at 18 inches.

Stand 13, Bur oak woods. Soil physical and chemical properties.

Property	Soil Depth, inches				
	0-6	6-12	12-18	18-24	36
Bulk density (g/cc)	1.052	-	1.621	-	-
Texture					
% Sand	52	59.5	63	57.5	50.5
% Silt	35	26	21	21.5	25.5
% Clay	13	14.5	16	21	24
Moisture properties					
15 Bar moist. (%)	18.91	10.70	6.51	7.37	6.45
1/3 Bar moist. (%)	26.59	18.24	13.57	15.16	16.44
Avail. moist. (%)	7.68	7.54	7.06	7.79	9.99
Organic matter (%)	9.24	4.97	2.58	3.07	2.86
pH	6.7	6.4	6.5	6.7	7.3
Nitrogen (%)	0.318	0.157	0.071	0.061	0.041
Phosphorus (ppm)	18.5	12.6	6.0	2.5	2.6
Potassium (ppm)	71.4	43.0	31.9	27.3	18.7
Calcium (ppm)	4220	2277	1670	3857	6233
Magnesium (ppm)	626	425	367	379	433
Sodium (ppm)	15.8	15.3	16.3	18.2	19.7
Hydrogen (ppm)	74.7	63.7	41.1	28.8	12.3
Base saturation (%)	78.1	70.7	73.4	81.0	90.9
Bases (meq/100g)	26.5	15.05	11.52	22.55	34.86
CEC (meq/100g)	33.91	21.38	15.60	25.41	36.08

Basis: Average of 3 replications, except bulk density with 6 at 6 inches and 3 at 18 inches.

Stand 14, Prairie. Soil physical and chemical properties.

Property	Soil Depth, inches				
	0-6	6-12	12-18	18-24	36
Bulk density (g/cc)	1.127	-	1.302	-	-
Texture					
% Sand	69	69.5	67.5	75.5	83
% Silt	20.5	13.5	13	10	9.5
% Clay	10.5	17	19.5	14.5	7.5
Moisture properties					
15 Bar moist. (%)	19.35	9.69	8.83	5.76	2.17
1/3 Bar moist. (%)	25.79	22.34	21.28	15.21	5.39
Avail. moist. (%)	6.44	12.65	12.45	9.45	3.22
Organic matter (%)	9.13	5.36	4.43	2.76	0.93
pH	7.8	8.0	8.0	8.0	8.1
Nitrogen (%)	0.368	0.174	0.127	0.076	0.030
Phosphorus (ppm)	1.1	1.6	1.4	1.4	0.75
Potassium (ppm)	12.5	6.9	7.8	8.1	6.7
Calcium (ppm)	9633	9513	9227	8753	6300
Magnesium (ppm)	928	568	420	323	177
Sodium (ppm)	16.7	17.4	18.1	16.6	23.3
Hydrogen (ppm)	0	0	0	0	0
Base saturation (%)	100	100	100	100	100
Bases (meq/100g)	55.90	51.33	49.69	46.51	33.08
CEC (meq/100g)	55.90	51.33	49.69	46.51	33.08

Basis: Average of 3 replications, except bulk density with 6 at 6 inches and 3 at 18 inches.

Stand 15, Bur oak savanna, prairie type. Soil physical and chemical properties.

Property	Soil Depth, inches							
	0-3	3-6	6-12	12-18	18-24	36	48	78*
Bulk density (g/cc)	-	1.558	-	1.615	-	-	-	-
Texture								
% Sand	93	95	95	96	96	96	96	95.5
% Silt	3	2	1.5	1	1	1	1	1
% Clay	4	3	3.5	3	3	3	3	3.5
Moisture properties								
15 Bar moist. (%)	3.59	2.37	1.78	1.33	1.17	1.21	1.09	0.96
1/3 Bar moist. (%)	4.21	2.73	2.05	1.62	1.53	1.48	1.45	1.53
Avail. moist. (%)	0.62	0.36	0.27	0.29	0.36	0.27	0.36	0.57
Organic matter (%)	1.60	0.96	0.71	0.49	0.43	0.39	0.38	0.30
pH	6.3	6.2	6.2	6.2	6.3	6.3	6.3	6.2
Nitrogen (%)	0.056	0.036	0.021	0.014	0.012	0.007	0.006	0.004
Phosphorus (ppm)	11.7	9.3	9.2	7.7	7.7	9.3	12.4	16.1
Potassium (ppm)	8.8	6.3	5.9	5.5	4.8	4.7	5.5	5.0
Calcium (ppm)	580	358	268	222	206	212	209	160
Magnesium (ppm)	79.0	48.7	39.6	36.1	31.6	29.2	27.3	29.6
Sodium (ppm)	0.2	0.9	0.4	0.8	0.0	4.0	5.8	0.2
Hydrogen (ppm)	24.7	20.1	15.6	11.9	7.4	12.3	7.8	20.6
Base saturation (%)	59.8	53.2	53.2	56.7	70.1	54.7	64.4	34.1
Bases (meq/100g)	3.57	2.21	1.69	1.42	1.30	1.33	1.31	1.06
CEC (meq/100g)	6.02	4.21	3.24	2.61	2.03	2.55	2.08	3.10

Basis: Average of 5 replications, except bulk density with 10 at 6 inches and 9 at 18 inches.

* 1 sample.

Stand 15, Bur oak savanna, juniper type. Soil physical and chemical properties.

Property	Soil Depth, inches						
	0-3	3-6	6-12	12-18	18-24	36	48
Bulk density (g/cc)	-	1.533	-	1.617	-	-	-
Texture							
% Sand	93	94	95	96	95.5	96	95.5
% Silt	3	2.5	1.5	1	1	1	1
% Clay	4	3.5	3.5	3	3.5	3	3.5
Moisture properties							
15 Bar moist. (%)	4.25	2.76	1.80	1.54	1.40	1.25	1.17
1/3 Bar moist. (%)	4.48	3.04	2.24	1.93	1.72	1.64	1.54
Avail. moist. (%)	0.23	0.28	0.44	0.39	0.32	0.39	0.37
Organic matter (%)	1.67	1.14	0.76	0.75	0.52	0.47	0.42
pH	6.1	5.8	5.8	5.7	5.7	5.9	5.9
Nitrogen (%)	0.055	0.040	0.022	0.017	0.012	0.010	0.007
Phosphorus (ppm)	10.6	10.9	10.3	9.7	9.8	12.5	14.4
Potassium (ppm)	10.6	5.8	4.36	4.0	3.7	3.9	4.2
Calcium (ppm)	554	323	241	206	196	195	191
Magnesium (ppm)	88.4	54.5	39.2	33.0	29.4	25.9	25.6
Sodium (ppm)	2.8	0.0	0.3	2.4	0.0	0.0	0.0
Hydrogen (ppm)	30.5	28.0	18.5	18.9	14.4	14.4	25.5
Base saturation (%)	54.6	44.4	48.3	44.7	49.1	49.6	37.9
Bases (meq/100g)	3.54	2.08	1.54	1.32	1.23	1.20	1.18
CEC (meq/100g)	6.56	4.85	3.38	3.20	2.66	2.63	3.70

Basis: Average of 5 replications, except bulk density with 11 at 6 inches and 9 at 18 inches.

Stand 15, Bur oak savanna, oak-prairie type. Soil physical and chemical properties.

Property	Soil Depth, inches						
	0-3	3-6	6-12	12-18	18-24	36	48
Bulk density (g/cc)	-	1.465	-	1.592	-	-	-
Texture							
% Sand	92.5	94	95	95.5	95.5	96	96
% Silt	3.5	2.5	1.5	1	1.5	1	1
% Clay	4	3.5	3.5	3.5	3	3	3
Moisture properties							
15 Bar moist. (%)	4.56	3.46	2.29	1.72	1.46	1.27	1.20
1/3 Bar moist. (%)	5.10	3.88	2.72	2.03	1.85	1.91	1.60
Avail. moist. (%)	0.54	0.42	0.43	0.31	0.39	0.64	0.40
Organic matter (%)	1.95	1.43	0.89	0.61	0.50	0.41	0.38
pH	5.8	5.9	5.8	5.8	5.7	5.8	6.0
Nitrogen (%)	0.064	0.047	0.028	0.016	0.011	0.008	0.005
Phosphorus (ppm)	21.8	22.0	20.4	20.1	21.2	24.8	18.0
Potassium (ppm)	10.5	7.1	8.4	6.4	6.0	6.0	6.2
Calcium (ppm)	619	511	302	212	184	171	169
Magnesium (ppm)	70.0	49.2	41.1	33.4	43.1	29.8	36.8
Sodium (ppm)	2.5	3.8	2.2	2.0	2.7	4.3	2.7
Hydrogen (ppm)	39.1	32.5	26.7	22.6	21.0	19.8	23.0
Base saturation (%)	48.7	46.1	40.9	37.9	38.0	38.1	38.1
Bases (meq/100g)	3.71	2.99	1.88	1.36	1.30	1.13	1.18
CEC (meq/100g)	7.58	6.22	4.53	3.60	3.38	3.09	3.46

Basis: Average of 5 replications, except bulk density with 10 at 6 inches and 10 at 18 inches.

Stand 15, Bur oak savanna, oak-juniper type. Soil physical and chemical properties.

Property	Soil Depth, inches						48*
	0-3	3-6	6-12	12-18	18-24	36	
Bulk density (g/cc)	-	1.515	-	1.637	-	-	-
Texture							
% Sand	92	94	95	96	96	95.5	96
% Silt	3.5	2.5	1.5	1	1	1	1
% Clay	4.5	3.5	3.5	3	3	3.5	3
Moisture properties							
15 Bar moist. (%)	5.29	2.97	2.10	1.58	1.52	1.52	1.44
1/3 Bar moist. (%)	6.05	3.47	2.46	1.92	1.82	2.04	1.94
Avail. moist. (%)	0.76	0.50	0.36	0.34	0.30	0.52	0.50
Organic matter (%)	2.27	1.23	0.80	0.56	0.46	0.46	0.43
pH	5.8	5.9	5.7	5.7	5.6	5.7	5.6
Nitrogen (%)	0.069	0.038	0.023	0.014	0.011	0.009	0.007
Phosphorus (ppm)	19.4	22.7	19.1	16.1	14.4	16.6	18.4
Potassium (ppm)	21.2	13.1	8.2	6.5	5.7	5.5	4.8
Calcium (ppm)	728	372	235	184	183	214	178
Magnesium (ppm)	95.0	60.4	41.9	30.2	27.2	34.0	31.7
Sodium (ppm)	0.1	0.2	0.5	0.0	0.0	0.7	0.4
Hydrogen (ppm)	32.9	31.28	23.8	18.5	17.7	18.5	16.0
Base saturation (%)	55.6	43.5	40.6	40.6	42.0	45.6	37.4
Bases (meq/100g)	4.48	2.39	1.54	1.18	1.15	1.37	1.17
CEC (meq/100g)	7.74	5.49	3.91	3.02	2.91	3.20	3.15

Basis: Average of 5 replications, except bulk density with 10 at 6 inches and 8 at 18 inches.

* 4 samples.

Stand 15, Bur oak savanna, aspen-oak type. Soil physical and chemical properties.

Property	Soil Depth, inches						
	0-3	3-6	6-12	12-18	18-24	36	48
Bulk density (g/cc)	-	1.298	-	1.554	-	-	-
Texture							
% Sand	85.5	91.5	94	94.5	93	94	94.5
% Silt	8.5	4.5	2.5	2	3	2	1.5
% Clay	6	4	3.5	3.5	4	4	4
Moisture properties							
15 Bar moist. (%)	12.58	5.30	2.93	2.12	1.68	1.48	1.56
1/3 Bar moist. (%)	14.79	6.68	3.64	2.53	2.34	2.33	2.41
Avail. moist. (%)	2.21	1.38	0.71	0.41	0.66	0.85	0.85
Organic matter (%)	6.13	2.72	1.21	0.76	0.50	0.41	0.42
pH	5.4	5.4	5.5	5.6	6.3	6.2	6.0
Nitrogen (%)	0.187	0.086	0.045	0.022	0.010	0.007	0.005
Phosphorus (ppm)	19.4	22.8	20.8	23.0	17.2	22.0	18.6
Potassium (ppm)	12.4	10.0	6.2	5.4	6.8	7.0	7.2
Calcium (ppm)	1264	518	281	230	255	222	219
Magnesium (ppm)	209.0	88.7	51.4	48.3	69.6	52.1	37.8
Sodium (ppm)	12.8	7.7	11.8	8.8	12.6	11.1	9.0
Hydrogen (ppm)	119.3	70.4	39.1	27.2	13.9	14.4	16.4
Base saturation (%)	41.9	36.4	33.5	35.0	54.5	55.3	48.9
Bases (meq/100g)	8.12	3.37	1.90	1.60	1.92	1.60	1.46
CEC (meq/100g)	19.96	10.36	5.77	4.30	3.30	3.03	3.10

Basis: Average of 4 replications, except bulk density with 8 at 6 inches and 6 at 18 inches.

Stand 16, Aspen-oak woods. Soil physical and chemical properties.

Property	Soil Depth, inches					
	0-4*	4-6*	0-6	6-12	12-18	18-24
Bulk density (g/cc)	-	-	1.309	-	1.450	-
Texture						
% Sand	57.5	64.5	55.5	44.5	39	38.5
% Silt	21.5	19.5	21	28.5	31	31.5
% Clay	21	16	23.5	27	30	30
Moisture properties						
15 Bar moist. (%)	11.00	8.09	11.44	8.44	8.35	7.32
1/3 Bar moist. (%)	19.41	13.56	19.26	18.94	19.29	17.52
Avail. moist. (%)	8.41	5.50	7.86	10.50	10.94	10.20
Organic matter (%)	3.48	2.66	3.46	4.30	4.85	4.11
pH	6.8	7.4	7.2	7.9	8.1	8.2
Nitrogen (%)	0.101	0.049	0.086	0.060	0.039	0.025
Phosphorus (ppm)	5.1	2.6	4.0	1.2	1.3	1.7
Potassium (ppm)	37.4	24.6	36.0	21.1	16.2	15.1
Calcium (ppm)	2760	3360	3295	8053	8467	8747
Magnesium (ppm)	710	938	905	779	756	903
Sodium (ppm)	11.6	17.2	13.1	12.7	15.9	15.3
Hydrogen (ppm)	47.3	0.0	11.8	0.0	0.0	0.0
Base saturation (%)	80.8	100	95.2	100	100	100
Bases (meq/100g)	19.78	24.65	24.06	46.78	48.66	51.26
CEC (meq/100g)	24.48	24.65	25.24	46.78	48.66	51.26

Basis: Average of 3 replications, except bulk density with 6 at 6 inches and 2 at 18 inches.

* 1 sample.

5.4 Plant Nutrient Concentration Data

Stand 2, Bur oak woods. Plant analyses.

Species	P(%)	K(%)	Ca(%)	Mg(%)	Na(%)	Al (ppm)
<i>Fraxinus pennsylvanica</i>	0.4224	1.6035	1.0809	0.2686	0	24.12
<i>Parthenocissus quinquefolia</i>	0.3636	1.7191	1.2634	0.3025	0	33.27
<i>Poa pratensis</i>	0.1864	1.9785	0.2428	0.1487	0	22.05
<i>Prunus virginiana</i>	0.3809	1.3818	3.3477	0.4238	0	55.63
<i>Quercus macrocarpa</i>	0.2037	1.0770	0.9460	0.2430	0	21.85
<i>Ribes cynosbati</i>	0.3714	3.0262	1.2061	0.2607	0	35.52
<i>Symphoricarpos occidentalis</i>	0.4190	2.2739	0.6131	0.3007	0	12.03
<i>Tilia americana</i>	0.2849	1.8678	1.4938	0.3682	0	31.38
<i>Ulmus americana</i>	0.2590	1.1875	1.4980	0.3302	0	31.38
<i>Xanthoxylum americanum</i>	0.4069	2.2044	1.3205	0.2488	0	28.68

Basis: One composite sample based on 2-6 selected individuals per species.

Stand 2, Bur oak woods. Plant analyses, cont.

Species	Fe(ppm)	Zn(ppm)	Cu(ppm)	Mo(ppm)	Mn(ppm)	B(ppm)
<i>Fraxinus pennsylvanica</i>	67.05	20.22	11.48	5.88	44.71	50.71
<i>Parthenocissus quinquefolia</i>	108.74	16.46	9.40	8.11	140.62	108.82
<i>Poa pratensis</i>	74.19	21.74	5.03	4.40	55.28	55.26
<i>Prunus virginiana</i>	82.29	13.17	6.61	7.66	402.54	91.44
<i>Quercus macrocarpa</i>	74.50	21.26	7.72	7.06	547.49	64.98
<i>Ribes cynosbati</i>	78.86	13.09	5.34	5.88	147.03	66.54
<i>Symphoricarpos occidentalis</i>	41.72	14.79	5.26	7.51	86.82	61.22
<i>Tilia americana</i>	76.52	19.38	9.08	7.81	128.31	79.17
<i>Ulmus americana</i>	118.92	17.00	6.53	6.77	170.97	50.39
<i>Xanthoxylum americanum</i>	68.45	21.55	9.16	6.47	97.13	74.21

Basis: One composite sample based on 2-6 selected individuals per species.

Stand 5, Bur oak savanna. Plant analyses.

Species	P(%)	K(%)	Ca(%)	Mg(%)	Na(%)	Al(ppm)
<i>Fraxinus pennsylvanica</i>	0.3022	1.2558	1.1410	0.2866	0	30.18
<i>Parthenocissus quinquefolia</i>	0.3480	1.1239	1.7818	0.3217	0	42.41
<i>Poa pratensis</i>	0.2331	1.8887	0.2642	0.1374	0	22.83
<i>Prunus virginiana</i>	0.3748	1.2558	2.7383	0.5019	0	58.13
<i>Quercus macrocarpa</i>	0.1985	0.9793	1.1709	0.2313	0	37.13
<i>Ribes cynosbati</i>	0.3558	3.1982	1.5495	0.2724	0	40.75
<i>Smilacina stellata</i>	0.3264	2.7730	0.7344	0.2269	0	51.76
<i>Symphoricarpos occidentalis</i>	0.3333	1.8613	0.6966	0.2636	0	16.03
<i>Ulmus americana</i>	0.3826	1.4577	1.5573	0.3245	0	44.21
<i>Xanthoxylum americanum</i>	0.3800	1.9881	1.3608	0.2225	0	46.02

Basis: One composite sample based on 2-6 selected individuals per species.

Stand 5, Bur oak savanna. Plant analyses, cont.

Species	Fe(ppm)	Zn(ppm)	Cu(ppm)	Mo(ppm)	Mn(ppm)	B(ppm)
<i>Fraxinus pennsylvanica</i>	74.34	15.05	8.84	5.88	60.58	62.03
<i>Parthenocissus quinquefolia</i>	131.72	15.23	9.00	7.06	169.59	67.69
<i>Poa pratensis</i>	78.39	17.45	5.34	2.79	35.50	55.67
<i>Prunus virginiana</i>	125.79	9.44	5.03	10.20	248.90	53.55
<i>Quercus macrocarpa</i>	100.82	24.48	9.08	5.88	342.86	84.83
<i>Ribes cynosbati</i>	87.30	15.32	4.00	6.47	161.74	54.94
<i>Smilacina stellata</i>	114.14	14.19	3.69	4.99	65.00	68.68
<i>Symphoricarpos occidentalis</i>	61.34	16.37	4.55	7.06	140.62	60.32
<i>Ulmus americana</i>	105.41	18.45	7.80	7.21	76.11	77.76
<i>Xanthoxylum americanum</i>	104.46	17.90	8.36	6.47	86.37	50.07

Basis: One composite sample based on 2-6 selected individuals per species.

Stand 6, Bur oak woods. Plant analyses.

Species	P(%)	K(%)	Ca(%)	Mg(%)	Na(%)	Al (ppm)
<i>Acer negundo</i>	0.3861	1.4181	2.2324	0.4435	0	180.08
<i>Corylus americana</i>	0.2072	0.5852	1.6067	0.3928	0	106.44
<i>Fraxinus pennsylvanica</i>	0.2840	1.2857	2.4102	0.3991	0	105.63
<i>Parthenocissus quinquefolia</i>	0.6573	1.8388	2.4870	0.3798	0	187.49
<i>Prunus virginiana</i>	0.3230	1.5234	2.7938	0.5247	0	137.89
<i>Quercus macrocarpa</i>	0.2884	0.9421	1.4366	0.2764	0	69.45
<i>Symphoricarpos occidentalis</i>	0.3740	1.4807	1.4083	0.4188	0	70.19
<i>Thalictrum dioicum</i>	0.4285	2.2739	0.9258	0.2445	0	92.33
<i>Xanthoxylum americanum</i>	0.6287	1.9881	1.5484	0.2748	0	94.17

Basis: One composite sample based on 2-6 selected individuals per species.

Stand 6, Bur oak woods. Plant analyses, cont.

Species	Fe(ppm)	Zn(ppm)	Cu(ppm)	Mo(ppm)	Mn(ppm)	B(ppm)
<i>Acer negundo</i>	310.36	19.94	5.74	11.71	105.68	106.36
<i>Corylus americana</i>	202.83	20.50	7.16	10.50	346.31	59.42
<i>Fraxinus pennsylvanica</i>	187.10	15.14	10.28	9.30	57.92	63.34
<i>Parthenocissus quinquefolia</i>	325.31	18.91	9.64	10.66	152.54	55.59
<i>Prunus virginiana</i>	214.33	11.43	4.16	11.56	247.95	79.67
<i>Quercus macrocarpa</i>	156.13	23.59	10.04	6.92	301.31	90.43
<i>Symphoricarpos occidentalis</i>	133.01	24.78	6.85	9.60	63.23	90.26
<i>Thalictrum dioicum</i>	183.30	44.62	14.23	7.36	48.22	47.88
<i>Xanthoxylum americanum</i>	182.15	21.36	10.36	7.81	85.03	47.72

Basis: One composite sample based on 2-6 selected individuals per species.

Stand 10, Aspen oak woods. Plant analyses.

Species	P(%)	K(%)	Ca(%)	Mg(%)	Na(%)	Al (ppm)
<i>Amelanchier humilis</i>	0.3748	1.5790	1.3549	0.4037	0	24.72
<i>Corylus americana</i>	0.1925	0.7483	1.3205	0.3615	0	50.37
<i>Crataegus rotundifolia</i>	0.2573	1.1491	1.3199	0.3648	0	40.31
<i>Elymus canadensis</i>	0.3463	2.1933	0.2332	0.1256	0	6.59
<i>Parthenocissus quinquefolia</i>	0.3861	1.5447	1.8803	0.3149	0	79.64
<i>Prunus virginiana</i>	0.3377	1.5888	2.8700	0.4704	0	61.13
<i>Quercus macrocarpa</i>	0.2305	0.9371	1.1445	0.2733	0	30.85
<i>Rosa blanda</i>	0.2892	1.2325	1.4855	0.3882	0	37.79
<i>Smilacina stellata</i>	0.2322	3.6689	0.8866	0.3245	0	41.41
<i>Symphoricarpos occidentalis</i>	0.3126	2.1442	0.8632	0.2944	0	17.25
<i>Viburnum rafinesquianum</i>	0.2503	1.7612	1.4071	0.4157	0	137.89

Basis: One composite sample based on 2-6 selected individuals per species.

Stand 10, Aspen oak woods. Plant analyses, cont.

Species	Fe(ppm)	Zn(ppm)	Cu(ppm)	Mo(ppm)	Mn(ppm)	B(ppm)
<i>Amelanchier humilis</i>	64.58	42.75	10.12	9.45	409.08	51.93
<i>Corylus americana</i>	91.85	19.38	11.48	9.60	418.15	48.29
<i>Crataegus rotundifolia</i>	89.02	34.84	9.32	7.51	60.13	52.17
<i>Elymus canadensis</i>	53.64	30.84	19.60	3.67	31.13	41.03
<i>Parthenocissus quinquefolia</i>	182.32	44.62	65.95	7.06	201.68	53.88
<i>Prunus virginiana</i>	113.34	18.27	18.70	9.60	322.25	65.23
<i>Quercus macrocarpa</i>	76.52	29.16	23.55	7.51	391.01	47.24
<i>Rosa blanda</i>	103.67	14.02	28.35	7.66	150.24	50.47
<i>Smilacina stellata</i>	80.88	14.71	18.37	7.06	53.07	52.74
<i>Symphoricarpos occidentalis</i>	58.41	23.10	9.80	6.62	135.15	51.61
<i>Viburnum rafinesquianum</i>	116.05	52.21	8.52	11.71	226.60	57.79

Basis: One composite sample based on 2-6 selected individuals per species.

Stand 11, Prairie. Plant analyses.

Species	P(%)	K(%)	Ca(%)	Mg(%)	Na(%)	Al (ppm)
<i>Amorpha canescens</i>	0.1096	0.7226	1.1755	0.2131	0	59.33
<i>Andropogon gerardi</i>	0.0872	1.0164	0.3634	0.1325	0	18.38
<i>Andropogon scoparius</i>	0.0863	0.9624	0.4175	0.1291	0	23.52
<i>Elymus canadensis</i>	0.0631	0.8743	0.2530	0.0637	0	42.19
<i>Heliopsis helianthoides</i>	0.1139	2.1902	2.2404	0.6130	0	121.73
<i>Liatris aspera</i>	0.0639	1.2857	1.2205	0.2315	0	48.30
<i>Petalostemum purpureum</i>	0.0760	0.9692	1.6922	0.1978	0	32.85
<i>Sorghastrum nutans</i>	0.0863	1.0080	0.4669	0.0916	0	34.23
<i>Veronicastrum virginicum</i>	0.0777	1.2558	0.7701	0.3590	0	63.31
<i>Zizia aptera</i>	0.1346	2.3289	2.0954	0.2845	0	125.88

Basis: One composite sample based on 2-6 selected individuals per species.

Stand 11, Prairie. Plant analyses, cont.

Species	Fe(ppm)	Zn(ppm)	Cu(ppm)	Mo(ppm)	Mn(ppm)	B(ppm)
<i>Amorpha canescens</i>	107.15	15.84	5.74	6.47	61.46	48.29
<i>Andropogon gerardi</i>	58.41	15.84	5.34	4.25	11.56	39.42
<i>Andropogon scoparius</i>	74.19	11.19	5.18	3.96	21.10	38.62
<i>Elymus canadensis</i>	98.77	15.58	2.19	4.99	23.71	46.35
<i>Heliopsis helianthoides</i>	161.18	12.67	6.61	12.62	29.38	142.40
<i>Liatris aspera</i>	71.08	41.94	4.79	5.43	39.88	48.29
<i>Petalostemum purpureum</i>	55.03	22.51	4.16	4.84	35.50	47.15
<i>Sorghastrum nutans</i>	85.26	18.18	2.82	4.11	36.81	39.02
<i>Veronicastrum virginicum</i>	103.35	13.25	8.60	8.85	20.23	42.08
<i>Zizia aptera</i>	151.58	46.40	5.82	8.26	53.51	46.19

Basis: One composite sample based on 2-6 selected individuals per species.

Stand 12, Bur oak woods. Plant analyses.

Species	P(%)	K(%)	Ca(%)	Mg(%)	Na(%)	Al (ppm)
<i>Acer negundo</i>	0.4813	1.6980	1.6790	0.3307	0	138.03
<i>Corylus americana</i>	0.2512	0.9726	2.1856	0.5828	0	144.38
<i>Elymus canadensis</i>	0.3697	2.7223	0.3998	0.1459	0	23.13
<i>Fraxinus pennsylvanica</i>	0.4077	1.4856	1.2391	0.3304	0	76.59
<i>Parthenocissus quinquefolia</i>	0.4155	1.9337	1.7198	0.4268	0	155.04
<i>Populus tremuloides</i>	0.2547	1.4790	1.7372	0.3191	0.0209	49.22
<i>Prunus virginiana</i>	0.4424	1.5921	2.4430	0.4276	0	77.98
<i>Quercus macrocarpa</i>	0.1899	0.0181	1.3106	0.2830	0	56.35
<i>Symphoricarpos occidentalis</i>	0.3506	2.4901	0.9320	0.3202	0	38.55
<i>Tilia americana</i>	0.3368	1.8983	2.3434	0.4669	0	80.53
<i>Uvularia grandiflora</i>	0.3005	2.4604	1.0161	0.4230	0	105.10
<i>Viburnum rafinesquianum</i>	0.2970	1.7012	1.2478	0.4982	0	141.97
<i>Xanthoxylum americanum</i>	0.4363	2.6065	1.3304	0.2125	0	85.29

Basis: One composite sample based on 2-6 selected individuals per species.

Stand 12, Bur oak woods. Plant analyses, cont.

Species	Fe(ppm)	Zn(ppm)	Cu(ppm)	Mo(ppm)	Mn(ppm)	B(ppm)
<i>Acer negundo</i>	261.74	19.94	7.24	8.41	103.42	72.55
<i>Corylus americana</i>	225.38	19.28	9.24	13.68	412.60	67.36
<i>Elymus canadensis</i>	67.36	24.48	6.61	4.40	33.75	50.96
<i>Fraxinus pennsylvanica</i>	148.01	19.28	26.78	7.51	64.12	47.96
<i>Parthenocissus quinquefolia</i>	250.18	13.93	10.76	10.50	118.33	60.81
<i>Populus tremuloides</i>	88.71	149.98	10.12	6.47	46.90	66.79
<i>Prunus virginiana</i>	126.43	10.39	4.87	8.70	108.83	87.17
<i>Quercus macrocarpa</i>	107.47	24.38	9.96	6.62	245.57	63.67
<i>Symphoricarpos occidentalis</i>	84.17	19.84	7.09	7.06	94.43	47.80
<i>Tilia americana</i>	147.20	17.72	11.08	8.41	60.58	72.55
<i>Uvularia grandiflora</i>	169.84	28.02	10.20	9.90	58.37	34.78
<i>Viburnum rafinesquianum</i>	204.16	39.99	7.48	12.01	117.42	49.09
<i>Xanthoxylum americanum</i>	151.58	22.61	14.87	6.32	64.12	54.04

Basis: One composite sample based on 2-6 selected individuals per species.

Stand 13, Bur oak woods. Plant analyses.

Species	P(%)	K(%)	Ca(%)	Mg(%)	Na(%)	Al(ppm)
<i>Acer negundo</i>	0.4112	1.4527	1.4761	0.2924	0	48.41
<i>Actaea rubra</i>	0.4025	2.8588	1.2148	0.2908	0	63.19
<i>Actaea rubra</i> (sample 2)	0.4155	3.2402	1.2646	0.3031	0	89.97
<i>Amelanchier sanguinea</i>	0.3480	1.1608	1.4171	0.3887	0	41.41
<i>Bromus inermis</i>	0.2832	2.4760	0.3030	0.1531	0	8.19
<i>Corylus americana</i>	0.2132	0.7705	1.3743	0.4533	0	41.08
<i>Crataegus rotundifolia</i>	0.2685	1.0669	1.5561	0.3300	0	38.44
<i>Parthenocissus quinquefolia</i>	0.4060	1.4692	1.7006	0.3277	0	74.70
<i>Populus tremuloides</i>	0.1925	0.8709	1.0883	0.2487	0.0167	29.50
<i>Prunus virginiana</i>	0.5021	1.3438	2.6735	0.5009	0	59.45
<i>Quercus macrocarpa</i>	0.2607	0.9167	0.8805	0.2281	0	28.99
<i>Ribes americanum</i>	0.4476	1.8565	1.9849	0.3488	0	66.00
<i>Rosa arkansana</i>	0.3325	1.3272	1.1588	0.3672	0	33.27
<i>Smilacina stellata</i>	0.3835	3.2462	0.8165	0.2817	0	55.16
<i>Symphoricarpos occidentalis</i>	0.4069	1.9449	0.7525	0.3261	0	16.50
<i>Viburnum rafinesquianum</i>	0.3186	1.7223	1.1158	0.4318	0	74.44

Basis: One composite sample based on 2-6 selected individuals per species.

Stand 13, Bur oak woods. Plant analyses, cont.

Species	Fe(ppm)	Zn(ppm)	Cu(ppm)	Mo(ppm)	Mn(ppm)	B(ppm)
<i>Acer negundo</i>	98.93	19.10	13.09	7.06	43.83	54.77
<i>Actaea rubra</i>	110.17	13.42	8.28	6.32	32.00	41.67
<i>Actaea rubra</i> (sample 2)	146.07	13.34	9.40	6.47	29.38	44.01
<i>Amelanchier sanguinea</i>	81.98	43.92	7.48	8.26	107.48	48.77
<i>Bromus inermis</i>	38.22	20.13	21.41	4.40	49.54	34.70
<i>Corylus americana</i>	82.13	24.18	15.93	10.20	148.41	62.44
<i>Crataegus rotundifolia</i>	73.41	26.38	6.37	7.51	19.80	30.07
<i>Parthenocissus quinquefolia</i>	180.01	15.23	10.92	7.36	88.16	47.07
<i>Populus tremuloides</i>	70.93	103.12	8.68	5.29	39.00	38.22
<i>Prunus virginiana</i>	103.83	15.23	7.48	9.75	120.14	72.14
<i>Quercus macrocarpa</i>	87.77	45.09	50.75	4.99	225.66	63.02
<i>Ribes americanum</i>	116.69	13.68	5.74	8.26	59.69	45.22
<i>Rosa arkansana</i>	86.36	13.00	8.84	7.81	54.83	56.16
<i>Smilacina stellata</i>	109.05	11.43	7.32	7.36	37.69	34.30
<i>Symphoricarpos occidentalis</i>	50.58	26.28	7.40	7.51	60.58	58.52
<i>Viburnum rafinesquianum</i>	131.56	58.83	9.00	9.75	128.77	58.19

Basis: One composite sample based on 2-6 selected individuals per species.

Stand 14, Prairie. Plant analyses.

Species	P(%)	K(%)	Ca(%)	Mg(%)	Na(%)	Al (ppm)
<i>Andropogon gerardi</i>	0.0838	0.6712	0.2296	0.3004	0	7.76
<i>Andropogon scoparius</i>	0.0803	0.6901	0.2657	0.3541	0	16.31
<i>Apocynum sibiricum</i>	0.1243	0.9557	1.1882	0.6206	0	24.52
<i>Liatris ligulistylis</i>	0.0424	0.6076	0.8805	0.5645	0	22.05
<i>Petalostemum purpureum</i>	0.0682	0.6437	2.0654	0.4310	0	28.47
<i>Salix humilis</i>	0.1226	0.6351	1.1095	0.4733	0	22.93
<i>Sorghastrum nutans</i>	0.0656	0.8557	0.1737	0.1875	0	12.65
<i>Thalictrum dasycarpum</i>	0.0708	0.4883	0.8387	0.5247	0	24.32
<i>Zizia aptera</i>	0.1277	1.2891	1.8979	0.7146	0	50.60

Basis: One composite sample based on 2-6 selected individuals per species.

Stand 14, Prairie. Plant analyses, cont.

Species	Fe(ppm)	Zn(ppm)	Cu(ppm)	Mo(ppm)	Mn(ppm)	B(ppm)
Andropogon gerardi	42.33	30.21	3.45	7.06	10.26	17.41
Andropogon scoparius	62.26	13.76	2.35	6.77	18.93	22.14
Apocynum sibiricum	61.49	17.81	6.85	12.62	53.95	48.29
Liatris ligulistylis	49.51	36.94	6.53	11.11	33.75	46.67
Petalostemum purpureum	35.19	18.18	4.00	8.26	28.51	43.12
Salix humilis	47.21	10.23	5.26	9.00	28.07	26.26
Sorgastrum nutans	55.64	27.20	1.49	4.99	17.19	21.04
Thalictrum dasycarpum	47.67	33.42	4.71	10.96	44.27	41.75
Zizia aptera	87.77	50.37	5.26	13.07	44.27	41.27

Basis: One composite sample based on 2-6 selected individuals per species.

Stand 15, Bur oak savanna. Plant analyses.

Species	P(%)	K(%)	Ca(%)	Mg(%)	Na(%)	Al (ppm)
<i>Amorpha canescens</i>	0.1925	0.8029	0.6417	0.1609	0	10.09
<i>Andropogon gerardi</i>	0.1864	1.2990	0.2703	0.1101	0	9.22
<i>Andropogon scoparius</i>	0.1605	0.9354	0.2525	0.0774	0	8.19
<i>Juniperus horizontalis</i>	0.1709	0.8489	1.0752	0.2586	0	72.19
<i>Liatris aspera</i>	0.1053	1.0703	1.0076	0.3470	0	23.43
<i>Populus tremuloides</i>	0.2020	0.8862	0.9528	0.2344	0.0336	18.19
<i>Quercus macrocarpa</i>	0.2132	0.8675	0.8287	0.1980	0	25.42
<i>Salix humilis</i>	0.1977	0.9506	0.8821	0.2121	0	26.03
<i>Smilacina stellata</i>	0.4121	2.4167	0.8978	0.2931	0	55.28

Basis: One composite sample based on 2-6 selected individuals per species.

Stand 15, Bur oak savanna. Plant analyses, cont.

Species	Fe(ppm)	Zn(ppm)	Cu(ppm)	Mo(ppm)	Mn(ppm)	B(ppm)
<i>Amorpha canescens</i>	61.80	25.28	9.00	4.25	92.64	36.14
<i>Andropogon gerardi</i>	56.10	26.28	4.87	2.93	17.63	14.19
<i>Andropogon scoparius</i>	45.23	18.18	3.92	1.91	20.67	20.17
<i>Juniperus horizontalis</i>	151.74	20.03	3.76	7.81	78.78	21.67
<i>Liatris aspera</i>	54.87	56.94	4.95	7.36	89.06	47.40
<i>Populus tremuloides</i>	56.72	188.02	8.36	5.73	72.11	46.99
<i>Quercus macrocarpa</i>	69.69	19.38	7.72	4.40	203.09	46.99
<i>Salix humilis</i>	63.65	104.44	6.85	3.81	53.95	27.13
<i>Smilacina stellata</i>	96.57	19.10	2.43	7.36	49.54	33.50

Basis: One composite sample based on 2-6 selected individuals per species.

Stand 16, Aspen-oak woods. Plant analyses.

Species	P(%)	K(%)	Ca(%)	Mg(%)	Na(%)	Al (ppm)
<i>Amelanchier alnifolia</i>	0.2443	1.2458	1.3047	0.5181	0	32.11
<i>Cornus racemosa</i>	0.2400	0.6832	1.9660	0.4726	0	38.44
<i>Corylus americana</i>	0.1968	0.5679	1.6025	0.4881	0	59.09
<i>Elymus canadensis</i>	0.1139	1.5970	0.2170	0.0986	0	5.84
<i>Maianthemum canadense</i>	0.2331	2.5709	0.9799	0.4627	0	52.70
<i>Populus tremuloides</i>	0.1623	1.0534	1.4761	0.3150	0	27.35
<i>Prunus virginiana</i>	0.1683	1.1591	2.8650	0.6351	0	57.41
<i>Quercus macrocarpa</i>	0.1838	0.7824	0.9855	0.2923	0	30.33
<i>Smilacina stellata</i>	0.1838	2.6375	1.2176	0.4800	0	48.87
<i>Thalictrum dasycarpum</i>	0.2210	1.6329	0.7679	0.3274	0	16.87
<i>Veronicastrum virginicum</i>	0.1519	2.0249	1.2466	0.5948	0	37.46
<i>Viburnum rafinesquianum</i>	0.1640	1.7093	1.3825	0.5364	0	79.77

Basis: One composite sample based on 2-6 selected individuals per species.

Stand 16, Aspen-oak woods. Plant analyses, cont.

Species	Fe(ppm)	Zn(ppm)	Cu(ppm)	Mo(ppm)	Mn(ppm)	B(ppm)
<i>Amelanchier alnifolia</i>	89.02	8.44	7.24	9.60	97.13	34.22
<i>Cornus racemosa</i>	78.86	8.67	5.10	9.00	35.94	21.35
<i>Corylus americana</i>	127.39	18.64	7.56	9.75	116.97	60.07
<i>Elymus canadensis</i>	37.77	7.03	3.61	1.76	19.36	13.96
<i>Maianthemum canadense</i>	110.80	11.35	6.21	10.05	37.25	35.90
<i>Populus tremuloides</i>	77.92	36.16	8.76	5.14	41.63	74.21
<i>Prunus virginiana</i>	106.68	7.99	6.05	11.56	52.19	60.89
<i>Quercus macrocarpa</i>	89.65	24.38	8.12	6.02	106.58	66.95
<i>Smilacina stellata</i>	107.15	6.66	5.66	9.45	24.58	36.70
<i>Thalictrum dasycarpum</i>	51.04	12.34	9.24	7.36	23.71	34.70
<i>Veronicastrum virginicum</i>	84.17	9.60	12.77	10.96	27.20	27.13
<i>Viburnum rafinesquianum</i>	152.23	15.05	6.77	10.20	61.90	75.53

Basis: One composite sample based on 2-6 selected individuals per species.

5.5 Instrument Construction

Details which are obvious from the accompanying photos and drawings (or which are of minor importance) are omitted in the discussion of individual instruments. The mention of brand names, companies, or trademarks does not constitute endorsement but is for informative or clarification purposes only.

5.51 Solar Radiometer

This instrument was constructed to measure the components of solar radiation. Total solar radiation is measured by orienting the instrument horizontally with the glass dome skyward (Fig. A-1); diffuse solar radiation is measured by employing the shade band (Fig. A-2); direct solar radiation is, then, the difference of these two measurements. Reflected solar radiation is measured by orienting the instrument horizontally with the glass dome groundward (Fig. A-3).

The instrument consists of three main components: the instrument housing, the glass dome, and the thermopile. The housing was milled from plexiglas (cast acrylic plastic) according to the dimensions shown in Fig. A-4. (The dimensions were determined by the availability of suitable glass domes.) Plexiglas was chosen because it is light in weight, has a low thermal conductivity, and it machines well. Although it has a high thermal expansion (compared to aluminum and other metals), within normal working temperatures (of the completed instrument) this expansion is negligible. Construction screws were 6/32 brass. Polished aluminum discs and tape (3M Scotch aluminum





Figure A-1. Photo of solar radiometer in position for sensing total solar radiation.




Figure A-2. Photo of solar radiometer with shade band in place for sensing diffuse solar radiation.




Figure A-3. Photo of solar radiometer in inverted position for sensing reflected solar radiation.




Figure A-4. Construction detail of the solar radiometer, side view. Compare the drawing with Figs. A-1, A-2, and A-3.




Figure A-4, cont. Construction detail of the solar radiometer, bottom view. Note the thermopile connection (soldered joints).

tape) (Figs. A-1 and A-4) were used as radiation shields, utilizing the very high reflectivity of polished aluminum (Reifsnyder and Lull, 1965). Housing surfaces not protected by polished aluminum were painted with 3M white Velvet Coating (Nextel, 101-A10), utilizing its high reflective properties. Aluminum rod was used for the instrument handle because of its high strength and light weight. As shown in Figs. A-1 and A-4, the handle was constructed in two parts, connected by a bolt, to aid in instrument orientation. A terminal block of plexiglas (because it has very low electrical conductivity) was provided as a convenience; 4/36 brass bolts were used as terminals. A level was attached to the instrument to provide ease in leveling.

The glass dome (2.9 inches in diameter) was purchased from Kahl Scientific Instrument Corporation (El Cajon, CA). The dome is ground and polished and is transparent to light in the 0.28 to 2.80 micron wavelength range.

The thermopile consists of 48 copper-constantan junctions, 24 each under the black and white surfaces (Fig. A-1). The spirally arranged junctions are connected in series as shown in Fig. A-5 (design form Szeicz, 1966). The thermopile support consists of a piece of phenolic 2.29 inches in diameter and 0.010 inches thick. The phenolic has a low thermal and electrical conductivity. Since the phenolic is manufactured with a high surface gloss, it was lightly sanded to remove the gloss and provide better paint adhesion.

Constantan wire (42 gauge), used in thermopile construction,




Figure A-5. Construction detail of the solar radiometer thermopile. Note the copper plating only extends partially through the thermopile support.

was woven through the thermopile support (Fig. A-5) and partially copper plated (the portion on the upper surface of the thermopile support) to provide the copper-constantan junctions. The copper was deposited electrolytically (Wilson and Epps, 1920) in a plating bath consisting of 200 g. of CuSO_4 , 40 g. H_2SO_4 (1N), and 4 l. H_2O (Szeicz, 1965). A $1\frac{1}{2}$ volt dry cell battery was used as the current source, with a resistance of about 550 ohms across the plating bath. A piece of 12 gauge solid copper wire placed in a plastic tube of slightly larger diameter was used as the source of copper (Szeicz, 1965); the amount of copper wire exposed to the plating solution within the tube determined the resistance across the plating bath. Prior to plating, the constantan wire on the underside of the thermopile support was covered with a clear fingernail polish (other high quality varnish may be used) at the holes in the thermopile support (Fig. A-5). (The varnish should extend about half way through the thermopile support, when it is of the proper consistency.) Then a piece of 32 gauge copper wire was woven through the remaining bare constantan wire on the underside of the thermopile support. This copper wire provided the connection to the current source. Since constantan wire has a very high resistance (in comparison with many other metals), a close coupling of the area to be plated with the current source was necessary. Next, the wire to be plated (upper surface of the thermopile) was cleaned with acetone and then carbon tetrachloride. (Acetone must be used sparingly to clean any stray fingernail polish from surfaces to

be plated, such as near the holes in the thermopile support.) Finally, in preparation for plating, the lower surface of the thermopile support was coated with paraffin to completely cover all constantan wire. The thermopile was then immersed in the plating bath for $1\frac{1}{2}$ hours. Following plating, the deposited copper was examined under 10-20X magnification for any anomalies or discontinuities in the plating. Extreme care must be taken in cleaning the wire to be plated; no air bubbles can be on the wire in the plating bath. (The latter may be avoided by slight agitation of the thermopile in the plating bath following immersion.) The paraffin was removed from the thermopile by immersion in three successive baths of boiling tertiary-butyl alcohol. The copper connection wire was then removed from the underside of the thermopile. The upper surface of the thermopile was spray painted with 3M Velvet Coating (two light, even coats) as shown in Fig. A-5 (black, Nextel, 101-C10; white, Nextel, 101-A10). Templates made of Herculene drawing film were used to aid in the painting. The thermopile output was tested prior to cementing into the radiometer housing. Thermopile lead wires were then soldered to the external lead. When the instrument was assembled, silicone seal (Dow Corning) was used to hold the glass dome in place. Prior to sealing the lower part of the instrument, a small bag of silica gel was enclosed to prevent any moisture condensation on the instrument interior. The lower cover (since it is removable) was sealed with a non-hardening silicone compound (Dow 3 compound), with the instrument in an

environment of low humidity (75°F, 31% r.h.).

The instrument was designed to incorporate the beneficial parts of several commercial instruments. Catalogs of instruments were consulted to prepare the basic design. Also, the articles by Monteith (1959a), Norris and Trickett (1968), and Szeicz, Monteith, and dos Santos (1964) were used in formulating ideas. The radial arrangement of black and white surfaces (Fig. A-1) was chosen over the concentric pattern of the Eppley design, even though the radial pattern is harder to construct, to provide equal areas of black and white absorbing surfaces (in the area of the junction).

The instruments were calibrated by comparing output readings with an Eppley pyranometer of known output. Readings were made on clear days and at times of low as well as high sun altitudes. Regressions of the instrument output on incoming radiation were calculated. Correlations of the readings ranged from 0.9995 to 0.9997 on 5 of the 8 radiometers constructed, and thus it was believed the straight line regression equations were adequate calibration equations for these instruments. On 3 instruments, which utilized glass domes with a slightly different shape, the correlations were 0.9963 to 0.9975. Fourth power polynomials were fitted to the output readings of these instruments to provide better calibration equations.

The shade band (Fig. A-2), used to provide measurements of diffuse solar radiation, was constructed of 0.020 inch aluminum, based on the design by Horowitz (1969). Similar

designs are also shown in Drummond (1964) and Robinson (1966). The basic differences in the Horowitz band and the one constructed, were that a) it was necessary for it to be light in weight, b) portable, and c) adapted to the previously described solar radiometers. The band has a width of 3.5 inches, slightly larger than the solar radiometer dome, and a diameter of 21.65 inches. The band was polished on the outside to provide high reflectivity and painted with the 3M black Velvet Coating on the inside to reduce reflection. A mounting frame was constructed of $3/8$ inch aluminum rod, $1/8 \times 3/4$ inch aluminum bar, and $1 \times 1 \times 1/16$ inch aluminum angle. Further explanation is left to Figs. A-6 and A-7.

There are other methods of constructing shade bands (Radiation Commission of IAM, 1958; Robinson, 1966; Robinson and Stock, 1964), but the one constructed appeared the simplest in construction and adjustments. The band must be adjusted for latitude and time of year, the former dictating the angle of the shade band above the horizontal plane of the radiometer, and the latter being a vertical displacement above the horizontal plane. The radiometer and shade band must be oriented towards true south to provide proper adjustment of the shade ring. Corrections may be applied for the portion of the sky blocked by the shade band (Radiation Commission of IAM, 1958; Robinson and Stock, 1964), but for practical purposes may be ignored.

5.52 Total Radiometer

This instrument was constructed to measure the components




Figure A-6. Construction detail of the solar radiometer shade band. (See also Fig. A-2.)




Figure A-7. Construction detail of the solar radiometer shade band frame. (See also Fig. A-2.)

of total radiation, both the downward and upward fluxes. The instrument is oriented horizontally above or below the surface from which a measurement is desired (Fig. A-8). The long-wave component of total radiation may be determined by the subtraction of the short-wave component, as measured with the solar radiometer, from the amount of total radiation. Net radiation is determined by the difference between the downward and upward fluxes of total radiation.

The instrument consists of 4 main components: the instrument housing, the polyethylene shielding domes, the interior thermocouple, and the thermopiles. The housing was milled from plexiglas (for the reasons explained in Sec. 5.51) according to the dimensions shown in Fig. A-9. (Dimensions were determined by the availability of suitable polyethylene domes.) Nylon screws (6/32) were used to connect the two radiometer halves (Fig. A-9). Polished aluminum tape and 3M white Velvet Coating were used for radiation shielding. An aluminum handle, plexiglas terminal block, (brass 4/36 terminal bolts) and level (Figs. A-8 and A-9) were used as explained in Sec. 5.51. Purge ports (Figs. A-8 and A-9) were provided as a means of circulating dry air through the instrument to prevent interior condensation, and to provide a means of inflating the polyethylene domes should they collapse (either from interior air contraction, differential air pressures, or from being touched). The ports consist of two 17 gauge stainless steel hypodermic needles (two in each half of the instrument) cut off so they extend through the side




Figure A-8. Photo of total radiometer in its sensing position.
Note the purge ports, terminal block, and the removable
polyethylene dome clamp ring.








Figure A-9. Construction detail of the total radiometer. Compare the drawing with Fig. A-8.



Bottom View (thermopile removed)



Side View showing screw hole detail



Cutaway Side View showing thermocouple and lead wires

Figure A-9, cont. Construction detail of the total radiometer. Compare the drawing with Figs. A-8 and A-9.

of the instrument (Figs. A-8 and A-9). A rubber inflator, a tube of CaSO_4 , and a purge port adapter (made from a disposable syringe) were used to provide the dry air source for purging.


Polyethylene domes (C. W. Thornthwaite Associates, Elmer, N. J.) 2.0 inches in diameter were used to shield the thermopiles. Polyethylene has a transparency in the 0.3 to 50 micron wavelength range (with minor exceptions) and therefore is a suitable material for shielding total radiometers (Fritschen, 1963, 1965b; Funk, 1959; Radiation Commission of IAM, 1958; Suomi and Kuhn, 1958; Tanner et al., 1969). Polyethylene cannot be cleaned easily, is easily damaged, and is subject to collapse (as mentioned above), and therefore the instrument was provided with a means of changing the domes. The domes are sealed with a non-hardening silicone seal (Dow 3 compound) and by means of a clamping ring and 8 nylon screws (6/32) (Figs. A-8 and A-9).

To measure the incoming and outgoing fluxes of total radiation, a means of differentiating the fluxes must be used. Normal net radiometers cannot do this. Thus, the total radiometers, as designed here, utilizes two net radiometers, both of which sense the difference in flux between the interior of the instrument and the outside environment. Then, the interior radiation flux is determined by the use of a thermocouple to sense the temperature (Fig. A-9) and by applying the Stefan-Boltzman Law (Kreith, 1965; Munn, 1966; Planck, 1914; et al.). The fluxes toward the instrument were then determined by subtraction.

The thermopiles were constructed from 42 gauge constantan wire, copper-plated to provide 32 junctions connected in series to provide 16 junctions sensing the thermal flux from each direction. Details of the thermopile are shown in Fig. A-10. The thermopile support was made from a piece of phenolic 0.010 inches thick and 1.89 inches in diameter. The constantan wire was woven radially and copper-plated as shown in Fig. A-10. A 32 gauge copper wire was woven under the constantan wire near the outside edge of the thermopile as a current source connection and to reduce the resistance to the area to be plated. Then the constantan wire was cleaned with carbon tetrachloride and the area not to be plated coated with paraffin. The rest of the plating procedure follows that described in the discussion of solar radiometers (Sec. 5.51).

The thermopiles, following plating, were tested, cemented into the radiometer housing, and the lead wires connected to the external leads. The interior of the instrument was painted with the 3M black Velvet Coating, as was the outward facing surface of the thermopile. Except for the rubber plugs of the purge ports and the removable polyethylene domes, the instrument was sealed with silicone seal (Dow Corning).

Calibration was accomplished by comparing output readings with those measured by a ventilated total hemispherical radiometer (Packard Bell) [similar to the instrument shown by MacKay, 1965]. Each half of the total radiometer was calibrated separately. The least squares method was used to fit third



Top View

Plating Detail



Figure A-10. Construction detail of the total radiometer thermopile (two used per instrument). Note the copper plating extends through the support disc to provide junctions on the upper and lower side of the support.

power polynomials to the data to provide calibration curves.

Output of the instruments are approximately 0.2 millivolts per $\text{cal cm}^{-2} \text{ min}^{-1}$ at an air temperature of about 21°C .

5.53 Soil Heat Flux Unit

This instrument (Fig. A-11) was designed to measure the heat flux to and from the soil. The direction of heat flow is determined by whether the output is positive or negative.

Philip (1961) discusses the necessary criteria of a heat flux unit. The designed instrument consists primarily of a non-shielded thermopile. A lead wire with junction block was provided as a convenience. Dimensions of the instrument are shown in Fig. A-12. Basically the thermopile consists of 42 gauge constantan wire copper-plated to form 30 junctions in series alternating between the upper and lower side of the unit. Plating was done as described in Secs. 5.51 and 5.52. The unit was semi-waterproofed using clear fingernail polish (a high quality varnish) and was painted with 3M black Velvet Coating. The design is similar to that of Deacon (1950); alternative (but similar) heat flux plates are described by Fuchs and Tanner (1968) and Hatfield and Wilkins (1950).

Calibration was accomplished by measuring the amount of heat flux through a box of fine sand (the same soil in which the instruments would be used) on a hot plate and by measuring the heat flux unit output. Regressions through the origin (Steel and Torrie, 1960) of heat flux on unit output provided the calibration constants. (Since zero heat flux requires zero



Figure A-11. Photo of the soil heat flux unit, with lead wire and junction block.




Figure A-12. Construction detail of the soil heat flux unit. Copper plating extends from top center to bottom center to provide junctions on both sides of the unit.

output from the heat flux unit, regressions through the origin were desired.) The equation $G = k \, dt/dz$ (adapted from Rose, 1966), where G ($\text{cal cm}^{-2} \text{ min}^{-1}$) is the heat flux through the soil, k ($\text{cal cm}^{-1} \text{ min}^{-1} \text{ }^{\circ}\text{C}^{-1}$) is the thermal conductivity, and dt is the difference in temperature ($^{\circ}\text{C}$) over the distance dz (cm), was used to determine actual soil heat flux. Thermometers of a known distance apart (with the heat flux unit between them) were used to measure the actual temperature of the sand. Based on the work by De Vries (1963), Kersten (1949), and Van Wijk (1963), and the moisture content of the sand, the thermal conductivity was determined to be $0.03345 \text{ cal cm}^{-1} \text{ min}^{-1} \text{ }^{\circ}\text{C}^{-1}$ (see also Smith, 1940; Smith and Byers, 1939). Kohnke and Nakshabandi (1964) and Van Wijk and Derksen (1966) discuss some of the factors which affect variability of thermal conductivity.

5.54 Instrument Tower

The instrument tower (Fig. A-13) was designed for the support of the previously described radiometers. The radiometer support arms were made of $\frac{1}{2}$ inch thinwall conduit, the cross arms from $\frac{1}{2}$ inch black pipe, and the uprights from $3/4$ inch black pipe. All the essential details can be seen in the photo (Fig. A-13). Cross tees and couplings were used to join the tower sections. The $\frac{1}{2}$ inch cross tees were reamed to accept the $\frac{1}{2}$ inch conduit, and were also drilled and tapped for a locking screw. The tower will also accept other instruments by use of appropriate fittings.



Figure A-13. Photo showing the constructed instrument tower.